MAY 07, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.sj2ecqe

Protocol Citation: Rengin Yamur Akbiyik, Sevval Uysalcan 2019. Protein Expression. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.sj2ecqe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 13, 2018

Last Modified: May 07, 2019

PROTOCOL integer ID:
 14682

Keywords: Protein Expression

Protein Expression

Rengin Yamur Akbiyik¹, Sevval Uysalcan¹

¹Istanbul Bilgi University

Rengin Yamur Akbiyik
 Istanbul Bilgi University

ABSTRACT

The protein expression is a process about synthesizing and modification for proteins in living organisms with using several laboratory techniques and devices. This technique is generally used for obtaining protein from the competent cell that is used for carrying plasmids that possess the protein that we want to express.

Protein expression involves two general steps that;

- Gene expression; which is continuum that converting DNA to the mRNA that can be translate into protein.
- Protein manufacturing; which is translation process.

MATERIALS

STEP MATERIALS

⊗ LB Broth Amresco Catalog #J106-2KG

Transform BL21 cells and plate them on Amp (Kan).

1

Afer overnight incubation, take colonies and transfer into 6..

2

🌡️ 37 °C

🧪 LB Broth Amresco Catalog #J106-2KG

🧪 6 mL LB

🧪 6 µL Ampicillin

When incubation process is done, transfer it to 1 L ZYM-505.

3

Centrifuge at 6000 rpm for 10 minutes.

4

Equipment

SORVALL LYNX 4000

NAME

Centrifuge

TYPE

Thermo Scientific

BRAND

75006580

SKU

🕒 00:10:00 Centrifuge 🌡️ 15 °C

Discard liquid and repeat centrifuge step for 3 times total.

5

00:10:00 2nd Centrifuge

00:10:00 3rd Centrifuge

Add Binding Buffer.

6

20 mL Binding Buffer

Transfer it to 50 ml falcon tube.

7

In this step 35 μ l of total sample should be obtained. Binding Buffer involves 5 ml Imidazole, 20 ml TBS, ddw up to 200 ml.

For the resuspension of pallet, tubes with Binding Buffer ca..

8

Sonication will be occurred.

9

For sonication, plastic test tubes are usually used for sonication. This sonication process is done at 3 cycles for 7 minutes. 00:07:00 Sonication for 1 cycle.

Repeat this step for 3 times total.

Centrifuge at 13000 rpm for 30 minutes.

10

00:30:00 Centrifuge 6 °C