SEP 26, 2019

Protein expression using *E. coli* strain BL21DE3

Robert Hooftman¹

¹Wageningen University

iGEM Wageningen 2019

Robert Hooftman
Wageningen University

ABSTRACT

In this protocol will be explained how to express proteins using *E. coli* strain BL21DE3 using a plasmid with a T7 promoter. The promoter of T7 polymerase from *E. coli* BL21DE3 has to be induced with IPTG.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.7i4hkgw

Protocol Citation: Robert Hooftman 2019. Protein expression using *E. coli* strain BL21DE3. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.7i4hkgw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Sep 21, 2019

Last Modified: Sep 26, 2019

PROTOCOL integer ID:
27964

Day 1

1 Prepare an overnight culture of the bacterium encoding for the protein you want to express.

16h

Day 2

- 2 Prepare an Erlenmeyer flask containing LB medium and the right antibiotics
- 3 Inoculate the Erlenmeyer flask 1:100 from your overnight culture and grow the bacteria at 37°C and 120 rpm.
- 4 Grow the culture to an OD600 of 0.6. This will usually take 2-3 hours. 3h
- 5 Take 1 ml of pre-induction sample and cold shock the culture for 30 minutes on ice. Keep the pre-induction sample in the fridge or freezer. 30m
- 6 Add IPTG to a final maximum concentration of 1 μM. IPTG is used because *E. coli*/BL21DE3 has its T7 polymerase gene under an IPTG sensitive promoter and we have used plasmids with a T7 promoter in our protein expression experiments
- 7 Grow the culture overnight (for 18hours) at 20°C and 120 rpm. 18h

Day 3

- 8 Measure the OD600 of a 10x diluted post-induction sample. Calculate the “real” optical density by multiplying this number by 10.

- 9 Take 1 ml of post-induction sample and spin this, together with the pre-induction sample down for 10 minutes at 4700 g, using a tabletop centrifuge. 10m
- 10 Resuspend the pellet of the pre-induction sample in 100µl of MQ. To correct for the difference in OD600 when you load the samples on a protein gel, resuspend the pellet of the post-induction sample in 100µl x
("real"OD600 post-induction sample)/(OD600pre-induction sample)
- 11 Prepare the samples to be loaded on a protein gel
- 12 Run both the pre-induction and post-induction sample on a protein gel