AUG 05, 2020

Protein Extraction

avinash.kale¹

¹domnic colvin

UM-DAE Centre for Excellence in Basic Sciences

avinash.kale

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bjdnki5e

Document Citation: avinash.kale 2020. Protein Extraction.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bjdnki5e>

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Aug 05, 2020

Last Modified: Aug 05, 2020

DOCUMENT integer ID:

40078

Protein Extraction

The four *Bacillus* bacteria were inoculated in 200 ml LB medium (Serva) and incubated at 37°C overnight. The cells were centrifuged at 13000 rpm for 15 minutes at 4°C to pellet the cells. The cell pellet was sonicated at amplitude 30%, pulse 10 seconds, gap 10 seconds, for a total time of 2 minutes: 30 seconds or 15 Cycles to obtain the cell lysate. The TRIzol extraction protocol was used with minor modifications(24). In brief, to the cell lysate, 1 mL of TRIzol reagent and 200 µL of chloroform were added. The solution was mixed vigorously and was centrifuged 14,000g for 15 minutes at 4°C. The Upper clear phase was carefully decanted and to the remaining phase, 300 µL of ethanol was added and incubated at room temperature for 3 minutes followed by centrifugation at 2000g for 15 minutes at 4 °C. To the supernatant, a 4-fold volume of ice-cold acetone was added and kept for overnight incubation

at -20°C. The protein pellet was washed thrice first with 0.3 M Guanidium-HCl in 95 % ethanol followed by ice-cold acetone two times. All the protein pellets were air dried and were subsequently dissolved in rehydration buffer (7 M urea, 2 M thiourea, 2 % CHAPS, Distilled water). Protein concentrations were estimated using Bradford assay.