

Aug 21, 2018 Version 2

## Protocol for *in vitro* transcription of DNA oligos by T7 polymerase V.2

DOI

[dx.doi.org/10.17504/protocols.io.sskeecw](https://doi.org/10.17504/protocols.io.sskeecw)



CGER<sup>1</sup>

<sup>1</sup>Audrey Lapinaite

The Center for Genome ...



Meredith Triplet

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.sskeecw](https://doi.org/10.17504/protocols.io.sskeecw)

**Protocol Citation:** CGER 2018. Protocol for *in vitro* transcription of DNA oligos by T7 polymerase. [protocols.io](#)  
<https://dx.doi.org/10.17504/protocols.io.sskeecw>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

We use this protocol and it's working

**Created:** August 21, 2018

**Last Modified:** August 21, 2018

**Protocol Integer ID:** 14892

### Abstract

This is a protocol for *in vitro* transcription of DNA oligos by T7 polymerase from the [Doudna Lab](#).

## Attachments



[RNA\\_in\\_vitro\\_transcr...](#)

226KB

## Guidelines

### Template design:

T7 polymerase requires a double-stranded promoter region for efficient template binding, but can then transcribe a single-stranded DNA template. Therefore, the following protocol uses an approach, whereby two oligos are annealed in order to generate a dsDNA promoter, followed by a single-stranded overhang that contains the reverse complement of the desired RNA sequence.

The reverse complement of the desired RNA sequence is ordered as a DNA oligo (assuming the length does not exceed ~80 nt), to which the reverse complement of the T7 promoter is appended at the 3' end Example template:

5'-TTTCTTAGCTGCCTATACGGCAGTGAA**CCTATAGTGAGT**

- **XXX** = reverse complement of T7 promoter
- **XXX** = reverse complement of the desired RNA seq
- **Note:** the consensus T7 promoter sequence contains the first 6 nucleotides that are transcribed. However, a dinucleotide is required for high efficiency transcription.

A separate, synthetic DNA oligo is ordered that contains the T7 promoter in the forward direction, which can be annealed with the above construct: T7oligo: 5'-TAATACGACTCACTATA-3'

**Note:** according to Ambion, for transcription of synthetic oligonucleotides, only the -17 to -1 positions of the promoter need to be double-stranded. This oligo takes this into account.

## Materials

### STEP MATERIALS

- ☒ 5x transcription buffer
- ☒ 1.5 nmol T7oligo
- ☒ 1.0 nmol template
- ☒ DEPC H<sub>2</sub>O
- ☒ 5N NaOH
- ☒ 1N NaOH
- ☒ RNase-free DNase **Promega**
- ☒ EtOH
- ☒ Silanization Solution I
- ☒ 15% urea polyacrylamide gel
- ☒ 40% Acrylamide-bisacrylamide (19:1)
- ☒ Urea
- ☒ Water
- ☒ 10% APS
- ☒ 1x TBE
- ☒ 1x TBE buffer
- ☒ RNase-free water
- ☒ Corning Spin-X filter tube
- ☒ 10x NEBuffer 3
- ☒ CIP (10 U/μl) **New England Biolabs**
- ☒ Phenol-chloroform solution
- ☒ 5x transcription buffer
- ☒ 1.5 nmol T7oligo
- ☒ 1.0 nmol template
- ☒ DEPC H<sub>2</sub>O
- ☒ 5N NaOH
- ☒ 1N NaOH
- ☒ RNase-free DNase **Promega**
- ☒ EtOH

- ☒ Silanization Solution I
- ☒ 15% urea polyacrylamide gel
- ☒ 40% Acrylamide-bisacrylamide (19:1)
- ☒ Urea
- ☒ Water
- ☒ 10% APS
- ☒ 1x TBE
- ☒ 1x TBE buffer
- ☒ RNase-free water
- ☒ Corning Spin-X filter tube
- ☒ 10x NEBuffer 3
- ☒ CIP (10 U/ $\mu$ l) New England Biolabs
- ☒ Phenol-chloroform solution

## Protocol materials

- ☒ Phenol-chloroform solution
- ☒ 1N NaOH
- ☒ 15% urea polyacrylamide gel
- ☒ Corning Spin-X filter tube
- ☒ 10x NEBuffer 3
- ☒ 5N NaOH
- ☒ 1N NaOH
- ☒ RNase-free DNase **Promega**
- ☒ 40% Acrylamide-bisacrylamide (19:1)
- ☒ Corning Spin-X filter tube
- ☒ EtOH
- ☒ 10% APS
- ☒ Urea
- ☒ 40% Acrylamide-bisacrylamide (19:1)
- ☒ 5N NaOH
- ☒ 15% urea polyacrylamide gel
- ☒ Water
- ☒ 10% APS
- ☒ 1.0 nmol template
- ☒ Phenol-chloroform solution
- ☒ 1x TBE buffer
- ☒ EtOH
- ☒ 1.5 nmol T7oligo
- ☒ 1x TBE buffer
- ☒ RNase-free water
- ☒ 5x transcription buffer
- ☒ 1x TBE
- ☒ DEPC H<sub>2</sub>O
- ☒ RNase-free water
- ☒ DEPC H<sub>2</sub>O

- ☒ Water
- ☒ Silanization Solution I
- ☒ 5x transcription buffer
- ☒ 10x NEBuffer 3
- ☒ 1.0 nmol template
- ☒ Silanization Solution I
- ☒ 1x TBE
- ☒ CIP (10 U/ $\mu$ l) New England Biolabs
- ☒ CIP (10 U/ $\mu$ l) New England Biolabs
- ☒ 1.5 nmol T7oligo
- ☒ RNase-free DNase Promega
- ☒ Urea
- ☒ 5x transcription buffer
- ☒ 1.5 nmol T7oligo
- ☒ 1.0 nmol template
- ☒ DEPC H<sub>2</sub>O
- ☒ 5N NaOH
- ☒ 1N NaOH
- ☒ RNase-free DNase Promega
- ☒ Silanization Solution I
- ☒ EtOH
- ☒ 15% urea polyacrylamide gel
- ☒ 40% Acrylamide-bisacrylamide (19:1)
- ☒ Urea
- ☒ Water
- ☒ 10% APS
- ☒ 1x TBE
- ☒ 1x TBE buffer
- ☒ RNase-free water
- ☒ Corning Spin-X filter tube
- ☒ 10x NEBuffer 3

 CIP (10 U/ $\mu$ l) New England Biolabs

 Phenol-chloroform solution

## Safety warnings

- ❗ Please refer to the SDS (Safety Data Sheet) for safety warnings and hazard information.

## Before start

### Stock solutions:

<b>5x transcription buffer:</b>	10 ml
150 mM Tris-Cl, pH 8.1	1.5 ml
125 mM MgCl <sub>2</sub>	1.25 ml
0.05% Triton X-100	50 µl
10 mM spermidine	100 µl
H <sub>2</sub> O	to 10mL
store at -20 °C	

<b>10x TBE</b>	1 L
Tris base	108 g
Boric acid	55 g
0.5 M Na <sub>2</sub> EDTA pH 8.0	40 ml
H <sub>2</sub> O	to 1 L

<b>0.5 M Na<sub>2</sub>EDTA</b>	0.5 L
Disodium ethylenediaminetetraacetate 2H <sub>2</sub> O	93 g
NaOH to get pH 8.0	10 g
H <sub>2</sub> O	to 500 ml
autoclave	

<b>10 % (w/v) APS</b>	50 ml
Ammonium persulfate	5 g
H <sub>2</sub> O	to 50 ml
store at -20 °C	

<b>1M spermidine</b>	
Spermidine	0.7 g
H <sub>2</sub> O	to 5 ml

store at -20 °C	
-----------------	--

<b>5N NaOH</b>	25 ml
NaOH	5 g
H <sub>2</sub> O	to 25 ml

<b>1N NaOH</b>	20 ml
5N NaOH	5 ml
H <sub>2</sub> O	to 20 ml

<b>10 % Triton X-100</b>	10 ml
Triton X-100	1 ml
H <sub>2</sub> O	9 ml
store at -20 °C	

<b>1% (w/v) bromophenol blue</b>	10 ml
bromophenol blue	100 mg
H <sub>2</sub> O	to 10 ml

<b>1% (w/v) xylene cyanol</b>	10 ml
xylene cyanol	100 mg
H <sub>2</sub> O	to 10 ml

<b>2x RNA Loading Dye</b>	30 ml
95% formamide	28.5 ml
1 mM EDTA	60 µl of 0.5 M
0.025% SDS	75 µl of 10%
0.025% bromophenol blue	750 µl of 1%
0.01% xylene cyanol	300 µl of 1%
store at -20 °C	

## Preparing 10 uM hybridized template

- 1 Mix the following in a final volume of 100  $\mu$ l:  
20  $\mu$ l 5x transcription buffer  
1.5 nmol T7oligo (15 uM final)  
1.0 nmol template (10 uM final)  
DEPC H<sub>2</sub>O to 100  $\mu$ l

 5x transcription buffer

 1.5 nmol T7oligo

 1.0 nmol template

 DEPC H<sub>2</sub>O

- 2 Heat at 70-80 °C for 2 minutes.

 70 °C Heating

 00:02:00 Heating

- 3 Slow cool on bench-top.

- 4 Use immediately or store at -20 °C.

### Note

For some applications, prior gel purification of the DNA template may be desirable. However, for my transcriptions, the oligos are not very long and I immediately gel purify the RNAs after transcription, so purity of the DNA oligo is not a major concern.

 -20 °C Storage

## Preparation of NTPs

- 5 Add 500  $\mu$ l of RNase-free H<sub>2</sub>O to the 70-80 mg of each NTP.

 500  $\mu$ L RNase-free H<sub>2</sub>O

- 6 Bring pH of each NTP to 6.8-7.2 by adding 5  $\mu$ l of 5N NaOH followed by 10-90  $\mu$ l of 1N NaOH. Monitor pH in the stripes with 6.5-10 range. Most adjustment will be needed for ATP, least - for UTP.

 5N NaOH

 1N NaOH

7

Measure absorbance of 500 fold dilution: 3 µl to 1.5 ml of H<sub>2</sub>O.

ATP  $\lambda_{\text{max}}$  259 nm,  $\epsilon_{\text{max}} 1.59 \times 10^4 \text{ cm}^{-1} \text{ M}^{-1}$

CTP  $\lambda_{\text{max}}$  271 nm,  $\epsilon_{\text{max}} 0.9 \times 10^4 \text{ cm}^{-1} \text{ M}^{-1}$

GTP  $\lambda_{\text{max}}$  253 nm,  $\epsilon_{\text{max}} 1.37 \times 10^4 \text{ cm}^{-1} \text{ M}^{-1}$

UTP  $\lambda_{\text{max}}$  259 nm,  $\epsilon_{\text{max}} 1.59 \times 10^4 \text{ cm}^{-1} \text{ M}^{-1}$

8 Calculate concentrations and required dilutions to reach 100 mM concentration each.

## Transcription reaction

9 Mix the following in a 100 µl reaction:

	<u>Final concentrations:</u>
20 µl 5X transcription buffer	(1X)
5 µl 100 mM ATP	(5 mM)
5 µl 100 mM CTP	(5 mM)
5 µl 100 mM GTP	(5 mM)
5 µl 100 mM UTP	(5 mM)
1 µl 1M DTT	(10 mM)
0.1 µl 1 mg/mL pyrophosphatase (Roche)	(1 µg/ml)
10 µl 1 mg/mL T7 polymerase	(100 µg/ml)
10 µl 10 µM hybridized template	(1 µM)
DEPC H <sub>2</sub> O to 100 µl	

## Note

My typical final yields on transcription reactions of this size are roughly 30 µl of 10-20 uM (300-600 pmol). For my biochemical purposes, this is more than enough material. However, the above reaction can easily be scaled up.

- ¶ 5 µL 100 mM ATP
- ¶ 5 µL 100 mM CTP
- ¶ 5 µL 100 mM GTP
- ¶ 5 µL 100 mM UTP
- ¶ 1 µL 1M DTT
- ¶ 0.1 µL 1 mg/mL pyrophosphatase (Roche)
- ¶ 10 µL 1 mg/mL T7 polymerase

10 Incubate at 37 °C, overnight.

⌚ 37 °C Incubation

⌚ 16:00:00 overnight incubation

11 DNase treat:

1. Add 5 units (5 µl) RNase-free DNase directly to reaction.
2. Incubate at 37 °C, 30 minutes.

⊗ RNase-free DNase Promega

¶ 5 µL RNase-free DNase

⌚ 37 °C Incubation

⌚ 00:30:00 Incubation

## Note

This step is optional and may be omitted.

12

Add equal volume (100 µl) gel loading buffer directly to transcription reaction.

¶ 100 µL Gel loading buffer

### Note

My RNAs run very close to the xylene cyanol dye. Therefore, I use a gel loading buffer that contains only bromophenol blue.

- 13 Gel-purify transcribed RNAs on a medium thickness, denaturing urea-polyacrylamide gel.

### Note

For my RNAs (~30 nt) I use a 15% gel. At this polyacrylamide concentration, with RNA from a 100  $\mu$ l transcription reaction, I get excellent separation of the desired RNA length from n+1 and n-1 contaminants (these arise from template-independent addition of nts by T7 polymerase; and synthetic template oligos that are not the full length). Optimization of the gel running time and acrylamide concentration will be necessary for gel-purifying RNAs of different length.


## Gel-purification

- 14

PAGE Fragment Resolution: Denaturing Conditions (6M Urea)

<b>% Acrylamide</b>	<b>Fragment Size</b>
<b>30</b>	<b>2-8</b>
<b>20</b>	<b>8-25</b>
<b>10</b>	<b>25-35</b>
<b>8</b>	<b>35-45</b>
<b>6</b>	<b>45-70</b>
<b>5</b>	<b>70-300</b>
<b>4</b>	<b>100-500</b>

Clean glass plates with EtOH and coat with 200 µl Silanization Solution I.



15 Assemble glass plates.

16 Prepare 15% urea polyacrylamide gel.

 15% urea polyacrylamide gel

- 17 Denaturing Urea Polyacrylamide Gel (300 ml for a gig gel: 35 × 25 cm)

	<b>10%</b>	<b>12.5%</b>	<b>15%</b>
10x TBE	30 ml	30 ml	30 ml
40% Acrylamide-bisacrylamide (19:1)	75 ml	93.75 ml	112.5 ml
Urea	108 g	108 g	108 g
Water	192 ml	173 ml	155 ml
Before casting add:			
10% APS	1.5 ml	1.5 ml	1.5 ml
TEMED	600 µl	600 µl	600 µl

After addition of APS and TEMED to Urea-Polyacrylamide mixture pour the gel immediately.

 40% Acrylamide-bisacrylamide (19:1) Urea Water 10% APS

- 18 Place comb and polymerize for 60 min.

 01:00:00 Polymerize

- 19 Assemble electrophoresis apparatus.

- 20 Remove comb.

21 Fill inner and outer buffer chambers with 1x TBE.

 1x TBE

22 Rinse wells with 1x TBE buffer.

 1x TBE buffer

23 Pre-run the gel for 30 min to heat-up and remove remaining urea. Set constant P=40 W, 3000 V, 400 mA.

 00:30:00 Pre-running gel

24 Load sample.

25 Run P=40 W, 3000 V, 400 mA for around 3 h, until the bromophenol blue band is migrating in the last ¼ of the gel.

 03:00:00 Running gel

26 Remove one of the glass plates and cover the gel with saran wrap. Flip. Remove another glass plate and cover all gel with saran wrap.

27 UV shadow band and excise with sterile razor and place in Falcon tube.

28 Break gel slices with p-1000 tip.

29 Crush and soak gel slices in equal volume of RNase-free water.

 RNase-free water

30 Elute RNA overnight by rocking at 4 °C.

 4 °C Eluting RNA

## Next day

31 Filter out gel pieces by spinning sample through Corning Spin-X filter tube. Centrifuge at top speed for 5 minutes, 4 °C.

 4 °C Centrifugation 00:05:00 Centrifugation

## Note

I cut the top off a P-1000 tip so that I can pipette as much as possible of the water-gel slurry into the Corning filter tube.

 Corning Spin-X filter tube

- 32 Add 30 µl 10x NEBuffer 3 to each sample, then 1 µl CIP (10 U/µl). Incubate at 37 °C for one hour.

 10x NEBuffer 3 30 µL 10x NEBuffer 3 CIP (10 U/µl) New England Biolabs 1 µL CIP 37 °C Incubation 01:00:00 Incubation

## Note

This step is only required if removal of the 5' triphosphate is required (e.g. if the RNA will subsequently be 5'-radiolabeled with T4 PNK.)

- 33 Phenol/chloroform extract with 300-500 µl phenol-chloroform solution, pH 8.0.

 Phenol-chloroform solution 300 µL Phenol-chloroform solution

- 34

Ethanol precipitate the RNA, by adding 1/10 volume 3 M NaOAc (pH 5.2), 1/100 volume 100x linearized acrylamide and 3 volumes 100% ethanol. [described in step 35-40]

below]

35 Incubate at -80 °C, at least 1 hour

 -80 °C Incubation

 01:00:00 Incubation

36 Spin down at top speed, 20 minutes, 4 °C.

 4 °C Spinning down

 00:20:00 Spinning down

37 Remove supernatant.

38

Wash pellet with 70% ethanol.

39

Spin down again and remove supernatant.

40 Dry pellets in speed vacuum, ~10 minutes.

 00:10:00 Drying pellets

41 Resuspend pellets in 30 µL DEPC H<sub>2</sub>O (or buffer) and store at -20 °C.

 300 µL DEPC H<sub>2</sub>O (or buffer)

 -20 °C Storage