AUG 11, 2020

Protocol for study of the in vitro effect of purified serum anti-anti-anti-SpA antibodies (Ab-3) on cultures of Staphylococcus aureus.

Angel A Justiz-Vaillant¹, Patrick E. Akpaka²

¹University of the West Indies St. Augustine;

²University of the West Indies. St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bjkekkte

Protocol Citation: Angel A Justiz-Vaillant, Patrick E. Akpaka 2020. Protocol for study of the in vitro effect of purified serum anti-anti-anti-SpA antibodies (Ab-3) on cultures of Staphylococcus aureus.. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjkekkte>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 11, 2020

Last Modified: Aug 11, 2020

PROTOCOL integer ID:
 40294

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

This test shows the ability of Ab-3 antibody to inhibit the growth of *S. aureus* in vitro. The amounts of samples tested were 18 samples from chicks fed with hyper-immune eggs and therefore have the Ab-3, and six samples from chicks fed non-hyperimmune eggs and definitively do not have the Ab-3. Inhibition of the growth of the bacteria plated out in blood agar was observed only when the Ab-3 was added to said culture media, suggesting that the anti-anti-anti-SpA (Ab-3) bound to the *S. aureus* and inhibited their ability to proliferate in culture. On the other hand, bacterial growth was observed when blanks or sera of chicks fed non-hyper-immune eggs were used because these samples lacked the protective Ab-3 [1].

Reference

Justiz Vaillant AA, Akpaka PE, Smikle M, McFarlane-Anderson N (2012) In vitro Inhibition of *Staphylococcus aureus* isolates by anti-antidiotypic antibodies to *Staphylococcal* protein (SpA). *J Vaccines Vaccin* 3:127. doi:10.4172/2157-7560.1000127.

MATERIALS

MATERIALS

⊗ Blood agar **Oxoid Microbiology Products - Thermo Fischer**

⊗ Culti-Loops™ *Staphylococcus aureus* subsp. *aureus* ATCC™ 33592™ **Thermo Fisher Catalog #R4607020**

SAFETY WARNINGS

- ! The risk associated with a particular microbe can be consulted in the following website of the CDC.
(<http://www.cdc.gov/biosafety/publications/BiologicalRiskAssessmentWorksheet.pdf>)

- 1 The investigation of the neutralizing ability of purified anti-SpA antibodies is carried out as follows: One ml of brain heart infusion (BHI) broths is placed in 10 sterile test tubes.
- 2 To each tube is added an equal volume of 10 µl of purified anti-anti-anti-SpA (Ab-3) antibodies in a concentration of 1.25-µg/µl.

- 3 All samples are tested in duplicate.
- 4 An inoculum of an ATCC *Staphylococcus aureus* strain (ATCC #33592) is prepared to 0.5 commercially prepared McFarland scale standard ($1=300 \times 10^6/\text{ml}$ bacteria concentration). Then, ten μl of this inoculum is added serially from tubes 1 to 10 (prepared for each sample) and the dilutions were reported in this study as a bacterial concentration of 0.1, 0.01, and 0.001.
- 5 The absorbance value of each tube at 600 nm is measured.
- 6 A sample from each tube is plated out on Blood agar and incubated overnight at 35°C .
- 7 Inhibition of the bacterial growth is observed in individual and pooled sera of chicks fed with anti-SpA hyperimmune eggs up to 28 days, 15 days, and eight days period.