FEB 28, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bbsfinbn

External link:
<https://doi.org/10.1371/journal.pone.0229682>

Protocol Citation: Shaohua Huang, Yi Xing, Lijuan Zhu, Ke Zhang 2020. Protocol for the systematic review of the efficacy of the fixed combination of latanoprost and timolol versus other fixed combinations for primary open-angle glaucoma and ocular hypertension.

protocols.io
<https://dx.doi.org/10.17504/protocols.io.bbsfinbn>

MANUSCRIPT CITATION:
 Xing Y, Zhu L, Zhang K, Huang S (2020) The efficacy of the fixed combination of latanoprost and timolol versus other fixed combinations for primary open-angle glaucoma and ocular hypertension: A systematic review and meta-analysis. PLoS ONE 15(2): e0229682. doi: [10.1371/journal.pone.0229682](https://doi.org/10.1371/journal.pone.0229682)

Protocol for the systematic review of the efficacy of the fixed combination of latanoprost and timolol versus other fixed combinations for primary open-angle glaucoma and ocular hypertension

PLOS One

Shaohua Huang¹, Yi Xing¹, Lijuan Zhu¹, Ke Zhang¹

¹Department of Ophthalmology, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, People's Republic of China

Shaohua Huang

ABSTRACT

Background: Fixed-combination (FC) therapy is used in primary open-angle glaucoma (POAG) and ocular hypertension (OHT) patients who require more than one medication to reach their target intraocular pressure (IOP). Currently, there are several FC therapies available for the treatment of glaucoma. The FC of latanoprost/timolol (LTFC) is a commonly used. We will conduct a systematic review to compare the IOP-lowering effects of LTFC with other FCs for patients with POAG and OHT.

Materials and methods: We will search PubMed, EMBASE, the Cochrane Library, and Web of Science for randomized-controlled clinical trials and cross-over studies. The outcomes are mean IOP and IOP fluctuation after one month of treatment. Meta-analysis will be carried out using RevMan (version 5.1) software. After conducting meta-analyses, we will rate the quality of each meta-analysis as high, moderate, low, or very low using the "GRADE" system.

ATTACHMENTS

[File Protocol.doc](#)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Jan 24, 2020

Last Modified: Feb 28, 2020

PROTOCOL integer ID:
32295

Keywords: primary open-angle glaucoma. ocular hypertension. latanoprost. timolol. meta-analysis