AUG 11, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bjk7kkzn

Protocol Citation: Angel A Justiz-Vaillant 2020. Protocol of preparation of a protein-LAG conjugated to horseradish peroxidase by the periodate method.. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bjk7kkzn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 11, 2020

Last Modified: Aug 11, 2020

PROTOCOL integer ID:
 40319

Protocol of preparation of a protein-LAG conjugated to horseradish peroxidase by the periodate method.

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

ABSTRACT

Peroxidase conjugated to proteins L, A and G (SpLAG-HRP) is a new development in the field of biochemistry. This versatile protein binds to more 500 species of animal IgGs. It makes it suitable for the study of zoonosis and the assessment of the antibody production to infectious microorganisms. This immunomarker is not available in a recombinant form as their counterpart SpLA-HRP, SpLG-HRP and SpAG-HRP. It can be used in ELISA, immunohistochemistry, Western blot analysis, Dot blot, and RIA [1,2].

References

1. Justiz-Vaillant AA, Akpaka PE, McFarlane-Anderson N, Smikle MF. Comparison of techniques of detecting immunoglobulin-binding protein reactivity to immunoglobulin produced by different avian and mammalian species. *West Indian Med J*. 2013;62(1):12-20.
2. Vaillant AJ, McFarlane-Anderson N, Wisdom B, Mohammed W, Vuma S, et al. (2013) Immunoglobulin-binding Bacterial Proteins (IBP) Conjugates and their Reactivity with Immunoglobulin in Enzyme-Linked Immunosorbent Assays (ELISA). *J Anal Bioanal Tech* 4: 175. doi:10.4172/2155-9872.1000175.

GUIDELINES

All reagents but specially the enzyme and the sodium periodate solution have to be prepared freshly before mixing it with the enzyme.

MATERIALS

MATERIALS

☒ Ammonium Sulfate **P212121**

☒ Sodium periodate **Bio Basic Inc. Catalog #SB0875.SIZE.100g**

☒ sodium borohydride **Sigma Aldrich Catalog #452882**

☒ Horseradish Peroxidase (HRP) type IV **Sigma Aldrich Catalog #P8375-25KU**

☒ Staphylococcal Protein-A **Sigma Aldrich**

☒ Protein-L from P. Magnus **Contributed by users**

☒ Streptococcal protein G by Sigma Aldrich **Contributed by users**

Pipettes

20ml to 1000 ml glass

Scale

Incubator

Refrigerator

Freezer

Centrifuges

- 1 Horseradish peroxidase (500 µg in 50 µl NaCO₃ , pH 9.6) is mixed with freshly made sodium periodate solution (1.71 mg/ml) followed by incubation in the dark for 2 h.
- 2 Mix 500 µg of staphylococcal protein-A (SpA) with an equal amount (500 micrograms) of a mix of horseradish peroxidase-sodium periodate.
Mix 500 µg of streptococcal protein-G (SpG) with an equal amount (500 micrograms) of a mix of horseradish peroxidase-sodium periodate.
Mix 500 µg of Peptostreptococcal protein-L (SpL) with an equal amount (500 micrograms) of a mix of horseradish peroxidase-sodium periodate.
- 3 The three mixtures are incubated separately for 3 hours at 4°C with gentle agitation.
- 4 Forty µl of freshly prepared NaBH₄ solution (5 mg NaBH₄ /ml 0.1 mM NaOH) is then added to each mixture and then, the preparations are centrifuged (13,000rpm., 10 minutes at RT). Add to each preparation cold saturated ammonium sulphate solution and centrifuge them again (10000rpm, 25 minutes at 4°C).
- 5 Mix the 3 preparations and incubate in the dark with gentle agitation.

- 6 The mixture is then centrifuged for 25 min at 4°C and recover the pellet at the bottom of the tube.
- 7 The pellet (SpLAG-HRP) is re-suspended in 500 µl of PBS pH=7.4 and dialysed against 1L of PBS for 24 h with 3 buffer changes.
- 8 An equal volume of glycerol is added to the dialysate followed by 200 µl of bovine serum albumin, BSA (20 mg/ ml).
- 9 The SpLAG-HRP conjugate is then stored at -20°C until further used.