


Jun 07, 2020

Protocol to generate Gastruloids (LSCB, EPFL)

 In 1 collection

DOI

dx.doi.org/10.17504/protocols.io.9j5h4q6



Stefano D Vianello¹, Mehmet Girgin², Giuliana Rossi², Matthias Lutolf³

¹Marine Research Station, Institute of Cellular and Organismic Biology (ICOB), Avademia Sinica;

²Laboratory of Stem Cell Bioengineering, Institute of Bioengineering, School of Life Sciences (SV) and School of Engineering (STI), Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland,;

³Laboratory of Stem Cell Bioengineering, Institute of Bioengineering, School of Life Sciences (SV) and School of Engineering (STI), Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland and Institute of Chemical Sciences and Engineering, School of Basic Science (SB), EPFL, Lausanne, Switzerland



Stefano D Vianello

Marine Research Station, Institute of Cellular and Organismi...

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.9j5h4q6>



Protocol Citation: Stefano D Vianello, Mehmet Girgin, Giuliana Rossi, Matthias Lutolf 2020. Protocol to generate Gastruloids (LSCB, EPFL). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.9j5h4q6>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: November 20, 2019

Last Modified: June 07, 2020

Protocol Integer ID: 30045

Keywords: gastruloids, mESCs, generating gastruloids from mouse embryonic stem cell, gastruloid generation protocol, generating gastruloid, mouse embryonic stem cell, embryonic stem cell, gastruloid, lutolf lab, cell, generation,

Abstract

Gastruloid generation protocol as performed in the Lutolf Lab, EPFL.

For previously published protocols, see:

Baillie-Johnson, Peter, et al. "Generation of aggregates of mouse embryonic stem cells that show symmetry breaking, polarization and emergent collective behaviour in vitro." *JoVE (Journal of Visualized Experiments)*105 (2015): e53252.

and

Mehmet, G., et al. "Generating Gastruloids from Mouse Embryonic Stem Cells." *Protocol Exchange* (2018).

Guidelines

This protocol assumes standard c



Materials

MATERIALS

✕ StemPro™ Accutase™ Cell Dissociation Reagent **Thermo Fisher Scientific Catalog #A1110501**

✕ PBS, pH 7.4 **Thermo Fisher Catalog #10010015**

✕ 50mL Reagent Reservoir White PS **VWR International (Avantor) Catalog #613-1184**

✕ 96-well Clear Round Bottom Ultra-Low Attachment Microplate **Corning Catalog #7007**

✕ GSK-3 Inhibitor XVI **Merck Millipore (EMD Millipore) Catalog #361559**

The following recipes were used to prepare the media used throughout the protocol:

10% Serum Medium (home made fresh):

to make 500mL

- 434mL DMEM, high glucose, with GlutaMAX™ [CAT# 61965059, Gibco™/Life Technologies]
- 50mL ES-grade Foetal Bovine Serum [CAT#16141-079, Gibco™/Life Technologies]
- 5mL Penicillin-Streptomycin 10000U/mL [CAT#15140122, Gibco™/Life Technologies]
- 5mL Non Essential Amino Acids 100X [CAT#11140035, Gibco™/Life Technologies]
- 5mL Sodium Pyruvate 100mM [CAT#11360039, Gibco™/Life Technologies]
- 1mL beta-mercaptoethanol 50mM [CAT#31350010, Gibco™/Life Technologies]

for final concentrations of:

- Penicillin: 100U/mL
- Streptomycin: 100ug/mL
- Non Essential Amino Acids: 0.1mM
- Sodium Pyruvate: 1mM
- Glutamax™ (= L-Alanyl-Glutamine, included in DMEM): 3.97mM
- beta-mercaptoethanol: 0.1mM

to make **10% Serum 2i/LIF Medium (made fresh)**, add:

- 3uM CHIR99021 (1:1000 from our 3mM stock,) [CAT#361559, Merck/Millipore]
- 1uM PD0325901 (1:500 from our stock) [CAT#S1036, Selleck Chemicals]
- 100u/mL LIF (1:1000 from our stock), [sourced in house]

N2B27 Medium (home made fresh):

to make 500mL

- 237mL Neurobasal™ Medium [CAT#21103049, Gibco™/Life Technologies]
- 237mL DMEM/F-12, with GlutaMAX™, [CAT#31331093, Gibco™/Life Technologies]
- 5mL Penicillin-Streptomycin 10000U/mL [CAT#15140122, Gibco™/Life Technologies]
- 5mL Non Essential Amino Acids 100X [CAT#11140035, Gibco™/Life Technologies]
- 5mL Sodium Pyruvate 100mM [CAT#11360039, Gibco™/Life Technologies]
- 2.5mL GlutaMAX™ Supplement, 200mM [= L-Alanyl-Glutamine, CAT#35050038, Gibco™/Life Technologies]
- 1mL beta-mercaptoethanol 50mM [CAT#31350010, Gibco™/Life Technologies]



- 5mL B27 Supplement, serum-free, 50X [CAT#17504001, Gibco™/Life Technologies]
- 2.5mL N-2 Supplement, 100X [CAT#17502001, Gibco™/Life Technologies]

for final concentrations of:

- Penicillin: 100U/mL
- Streptomycin: 100ug/mL
- Non Essential Amino Acids: 0.1mM
- Sodium Pyruvate: 1mM
- GlutamaMAX™ (= L-Alanyl-Glutamine, also included in DMEM/F-12): 1mM + 2.50mM = 3.50mM
- beta-mercaptoethanol: 0.1mM


Troubleshooting

Before start

Starting grounds: a healthy culture of adherent mouse Embryonic Stem Cells (mESCs), grown in DMEM-10%Serum, +2i +LIF, split every two or three days. Gastruloid generation is done at the time of splitting, and simultaneously with it.


















Preparation of the cell suspension

- 1 Using a vacuum line+glass pasteur pipette, aspirate out the culture medium and replace with  3 mL PBS-/- , for a short wash

Note

When removing liquid during washes, do not completely dry out the cells. The surface of the well should still look glossy.


- 2 Aspirate out the PBS, and replace with  500 μ L Accutase.
- 3 Let Accutase act for ~  00:03:00 ,  Room temperature , tapping the sides of the plate to ease dissociation, and until most cells are floating (as clumps or single cells)
- 4 Slightly tilt the plate, and use a P1000 to pipette the suspension up and down to further break down cell clumps. Use each ejection to wash the surface of the plate, so to collect as many cells as possible.
- 5 Transfer the Accutase-cell suspension to a clean 15mL Falcon tube labelled "cells"
- 6 Use ~  4.5 mL DMEM-10%Serum to further wash the surface of the well, again, pipetting up and down and hoping to collect any previously missed cells. Transfer this secondary cell suspension in the same tube as before (total volume  5 mL)
- 7 Centrifuge  200 x g ( 1000 rpm , (15.5 cm rotor radius))  00:04:00 min
 4 °C
- 8 Using a vacuum line+glass pasteur pipette, aspirate the supernatant without disturbing the pellet, redissolve the pellet in  10 mL PBS-/-, and centrifuge again  200 x g
( 1000 rpm , (15.5 cm rotor radius))  00:04:00 min  4 °C
- 9 Repeat  [go to step #8](#) for a second PBS-/- wash

- 10 Using a vacuum line+glass pasteur pipette, aspirate the supernatant without disturbing the pellet, and resuspend the pellet in  1 mL N2B27



Safety information





It is important to thoroughly resuspend the pellet at this point, as a single cell suspension is needed for accurate counting later on. Resuspend the pellet by pipetting several times (~20) with a P1000, then switching to a P200, and then to a P20

Cell counting and Gastruloid seeding

- 11 Load  10 μ L of the N2B27-cell suspension into a manual cell counter (Neubauer chamber/haemocytometer), and calculate the concentration of cells in your suspension.


Note

Each well of the 96well plate will receive one  40 μ L drop of N2B27. Each drop should contain 300 cells, i.e. a concentration of 300 cells /  40 μ L = 7.5 cells/ μ L



To fill an entire 96well plate with  40 μ L drops,  40 μ L x 96 well =  3.840 mL N2B27 are needed. For simplicity,  5 mL N2B27 suspension are prepared: at 7.5 cells/ μ L, this requires 37500 cells.

Note

At this point, cells can also be used to seed a new culture and keep propagating the line. We plate from 65000 to 75000 cells in a well of a tissue-culture-treated 6well plate, filled with 2mL prewarmed DMEM/10%Serum+2i/LIF

- 12 To  5 mL fresh N2B27, add the volume of cell suspension carrying 37500 cells, as calculated before. (If you want to prepare more than one plate of gastruloids, scale the volume of N2B27 and of cells accordingly)




- 13 Load the  5 mL N2B27-cell suspension into a multichannel pipette reservoir, and dispense  40 μ L of this solution to each well of an ultra-low-adhesion 96well plate

Safety information

Make sure that the cell suspension in the reservoir is always well mixed, to ensure homogeneous dispensing in the wells. At each transfer, pipette up and down several time with the multichannel, and slightly agitate the reservoir from time to time.

- 14 Lightly tap the plate against the surface of the hood to make sure all drops are at the very bottom of each well, doublecheck under the microscope for the presence of cells in the drops, and leave the plate in a humidified incubator, 5%CO₂, undisturbed



 24:00:00 h

This is considered the beginning of Day 1 (D1), t=0h-24h


Growing Gastruloids

- 15 **Beginning of Day 2 (D2), t=24h-48h**

No intervention needed at this timepoint. The Gastruloids will keep developing in their






 40 μ L of N2B27. Leave to grow  24:00:00 h more

Expected result


Gastruloids should look like a small ( 100 μ m) cluster, still in the process of aggregating. Individual cells might still be visible in the surroundings

- 16 **Beginning of Day 3 (D3), t=48h-72h**






Prepare  16 mL N2B27, adding CHIR99021 to a final concentration of  3 micromolar (μM) . Load this into a multichannel pipette reservoir, and dispense  150 μL of this solution to each well of the plate (i.e. total volume of  190 μL per well). Place back in the incubator for  24:00:00 h


Expected result

Gastruloids should look like a small, clean, translucent sphere of around  $\pm 200 \mu\text{m}$ diameter. No additional (satellite) spheres should be seen around it.




17 Beginning of Day 4 (D4), t=72h-96h

Using a multichannel pipette, remove  150 μL of the medium from each well and replace with  150 μL fresh N2B27. Place back in the incubator for  24:00:00 h

Expected result

Gastruloids are round spheres of around  $\pm 300 \mu\text{m}$ in diameter. Extensive cell shedding is expected as a consequence of the CHIR pulse.

18 Beginning of Day 5 (D5), t=96h-120

Using a multichannel pipette, remove  150 μL of the medium from each well and replace with  150 μL fresh N2B27. Place back in the incubator for  24:00:00 h

Expected result

Gastruloids morphology is no longer symmetrical: they look like a ovoid with length of around $\pm 600\ \mu\text{m}$. The protruding part is more translucent than the rest, and this is the part that will elongate in the next 24h.

At $t=120\text{h}$, the Gastruloid looks like a bowling pin with length of around $\pm 700\ \mu\text{m}$ - $\pm 800\ \mu\text{m}$. A thinner extension protrudes at the "posterior", more translucent than the denser anterior (spherical).

Note

Day 6 (D6) and Day 7 (D7), $t=144\text{h}$, 168h

Culture can be extended up to 168h (i.e. 7 days total). This can be done by continuing the daily medium changes as above, or by transferring individual Gastruloids to separate wells of a 24well plate, in $\pm 800\ \mu\text{L}$ fresh N2B27, and replacing half of the medium one day later. The plate is kept shaking on an orbital shaker, in the incubator for both additional days. If keeping the Gastruloids in the original 96well plate, no shaking is required but there is an increased risk of the Gastruloids adhering to the sides of the well and degenerating.