JUN 07, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.7tzhnp6

Protocol Citation: Stefano D Vianello, Mehmet Girgin, Giuliana Rossi, Matthias Lutolf 2020. Protocol to immunostain Gastruloids (LSCB, EPFL). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.7tzhnp6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 01, 2019

Last Modified: Jun 07, 2020

PROTOCOL integer ID:
 28249

Keywords: Gastruloids, Immunostaining, Immunofluorescence

🌐 Protocol to immunostain Gastruloids (LSCB, EPFL)

Stefano D Vianello¹, Mehmet Girgin², Giuliana Rossi², Matthias Lutolf³

¹Marine Research Station, Institute of Cellular and Organismic Biology (ICOB), Avademia Sinica;

²Laboratory of Stem Cell Bioengineering, Institute of Bioengineering, School of Life Sciences (SV) and School of Engineering (STI), Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland;

³Laboratory of Stem Cell Bioengineering, Institute of Bioengineering, School of Life Sciences (SV) and School of Engineering (STI), Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland, and Institute of Chemical Sciences and Engineering, School of Basic Science (SB), EPFL, Lausanne, Switzerland

Stefano D Vianello

Marine Research Station, Institute of Cellular and Organismi...

ABSTRACT

Standard protocol used for immunostaining Gastruloids (aggregates of mouse embryonic stem cells) in the Lutolf Lab, EPFL.

For a published protocol on how to generate Gastruloids, see:

CITATION

Beccari et al. (2018). Generating Gastruloids from Mouse Embryonic Stem Cells. Protocol Exchange.

LINK

[10.1038/protex.2018.094](https://doi.org/10.1038/protex.2018.094)

GUIDELINES

All handling and transfer of Gastruloids from one solution to the next is done with a *cut* P1000 tip (to avoid damage). The tips should be coated in BSA- or serum-containing solution (in our case, the blocking PBS+FT solution itself). It is very important that all tips are coated, as Gastruloid will otherwise stick to the plastic and will be lost.

To transfer Gastruloids with minimal carry-over of liquid from one solution to the other, keep the pipette vertical until they all accumulate to the bottom of the tip. Slightly push to create a small hanging drop, and touch the surface of the new solution. All Gastruloids will transfer from the tip to the new solution. Hence, when the protocol calls e.g. for three PBS washes, three wells of a 6 well plate are filled with PBS, and Gastruloids moved serially from one to the other.

MATERIALS

MATERIALS

⊗ DAPI **Thermo Fisher Scientific Catalog #D1306**

⊗ Triton X-100 **Sigma Aldrich Catalog #T8787**

⊗ 4% Paraformaldehyde in PBS **Alfa Aesar Catalog #J61899-AK**

⊗ Permount mounting medium **Fisher Scientific**

⊗ Thermo Scientific™ SuperFrost Plus™ Adhesion slides **Thermo Fisher Scientific Catalog #J1800AMNZ**

⊗ PBS, pH 7.4 **Thermo Fisher Catalog #10010015**

⊗ Embryonic stem-cell FBS, qualified, US origin **Thermo Fisher Catalog #16141079**

⊗ Fluoromount-G **Southern Biotech Catalog #0100-01**

⊗ Greiner CELLSTAR® 6 well culture plates **greiner bio-one Catalog #657185**

⊗ Falcon™ 48well Polystyrene Microplates **Falcon Catalog #353078**

⊗ #1 Micro Cover Glass 22mmx22mm **Electron Microscopy Sciences Catalog #72200-10**

The following recipes were used:

PBS-FT (PBS+FBS+ 0.2% Triton)

to make 500mL

- 450mL PBS, 1X [CAT#10010015, Gibco™/Thermo Fisher Scientific]
- 50mL ES-grade FBS [CAT#16141079, Gibco™/Thermo Fisher Scientific]
NOTE: could be replaced with BSA, less expensive
- 0.1mL Triton-X100 [CAT#T8787, Sigma Aldrich]

4% PFA in PBS (if not buying pre-prepared one):

NOTE: work under a hood, wearing a N95 mask, and appropriate protective equipment

to make 500mL

- Fill glass beaker with 400mL PBS, 1X [CAT#10010015, Gibco™/Thermo Fisher Scientific]
- Add magnetic stirrer
- Add 20g granulated PFA powder [CAT#0964.1, Carl Roth]
- Heat to 60°C (NOT HIGHER!) on a stirring hotplate, the solution will become transparent from 55°C to 60°C
- When powder is dissolved, top up with PBS to 500mL total volume
- Adjust to pH7.4 with HCl or NaOH (5 drops 1MHCl)

Aliquot and store at -20°C medium- to long term

SAFETY WARNINGS

- ⚠ This protocol involves the use of 4% Paraformaldehyde (4%PFA). Perform all steps involving 4%PFA under a chemical hood, and wear appropriate body and eye protection. Do not inhale vapours.

BEFORE START INSTRUCTIONS

- Prepare PBS+FT (cfr. "Text Materials" for recipe)
- Coat wells of a 6-well plate with serum-containing buffer to prevent Gastruloids from sticking to it. Leave~30min at room temperature
- Bring 4%PFA to Room temperature

Fixing and primary antibody stain (D1)

1 Harvesting Gastruloids:

- 1.1** Using a cut P1000 tip to avoid accidental damage (cfr. "Guidelines"), collect Gastruloids from each well of the 96-well plate by placing the tip straight down to the bottom of each well, and aspirating up. Move to the next well until full tip capacity is reached.

Note

To increase reproducibility of analysis, do not collect Gastruloids from the outer wells of the plate (all 36 border wells). These tend to develop differently from all other Gastruloids, probably due to evaporation of the medium. We only collect Gastruloids from the inner 60 wells of the plate.

- 1.2** Deposit collected Gastruloids in a 15mL centrifuge tube as you go
- 1.3** When collected all Gastruloids needed, wait for them to sediment to the bottom of the tube, and remove as much medium as possible (vacuum pump+glass pipette)

1.4 Dilute away left-over traces of medium by gently resuspending the Gastruloids in ~5mL PBS-/-

2 Fixing (4%PFA):

2.1 Remove the serum-containing medium (in our case, PBSFT) from the 6-well plate (can keep and recycle to coat tips; cfr. "Guidelines") and replace with 2mL 4%PFA

Safety information

PFA is toxic. Operate under a fume hood and wear appropriate hand, body, and eye protection.

2.2 Collect all Gastruloids from the tube, wait for them to sink at the bottom of the tip, and touch the surface of the PFA vertically to transfer them with minimal liquid transfer (cfr. "Guidelines")

Safety information

Coat a cut P1000 tips in PBS+FT to avoid Gastruloids sticking to it!

Note

48h Gastruloids are particularly tricky to work with: they will stick and clump together, and they are almost invisible to the naked eye. To facilitate working with this timepoint, just push the whole contents of the tip into the next wash, without waiting for them to sink to the bottom of the tip, and without minding the carry over of medium from the previous wash. If possible, work with an excess of Gastruloids to compensate for loss during staining and washes.

2.3 Cover with aluminium foil to reduce fluorescence loss (when using reporter lines)

2.4 Incubate at 4 °C 02:00:00 , with or without orbital shaker

3 Washes (to remove PFA):

3.1 Fill wells of a 6-well plate with 4mL PBS-/- and transfer fixed Gastruloids using the same technique as above

3.2 Cover with aluminium foil and wait 00:10:00 min

3.3 repeat for two more washes [go to step #3.1](#)

II

Note

If short on time, it is not necessary to wait 10 minutes for each wash. Gastruloids can be transferred serially across three PBS-filled wells, and left 10 minutes only in the last one.

PAUSE POINT: you can leave Gastruloids in the last PBS wash, for months, 4 °C , protected from light

4 Blocking:

4.1 Transfer Gastruloids to a well filled with 2mL PBS+FT

4.2 Block for 01:00:00 h, Room temperature , with or without shaker

5 Primary antibody staining:

5.1 For every antibody combination and sample condition you want to stain, prepare a 1.5mL Eppendorf tube with 500uL of the appropriate antibodies in PBSFT. Include DAPI 1:500.

5.2 Transfer each antibody solution in a separate well of a 48 well plate, and transfer equal number of Gastruloids to each well

5.3 Cover in aluminium and incubate 24:00:00 , 4 °C , on a orbital shaker

Secondary antibody stain (D2)

6 Washes (to remove 1ry Ab):

6.1 Wash Gastruloids by transferring them to a well of a 6-well plate filled with 3mL PBS+FT

Note

Previous versions of this protocol call for PBS-/- instead of PBS-FT. There is a much higher increase of Gastruloids still sticking to the bottom of the wells in this case however. PBS-FT seems to be the safest option.

Safety information

You will need a separate PBS-FT well for each different sample condition, do not mix all Gastruloids into the same well

6.2 Wait for 00:20:00 min

6.3 Repeat for two more washes [go to step #6.1](#) (total: 01:00:00 h)

Note

Again, if short on time, it is not necessary to wait 20 minutes for each wash. Gastruloids can be transferred serially across three PBSFT-filled wells, and left 20 minutes only in the last one.

7 Secondary antibody staining:

7.1 For every antibody combination and sample condition stained, prepare 500uL of the appropriate secondary antibodies in PBSFT. Include DAPI 1:500 at this step too.

7.2 Transfer each antibody solution in a separate well of a 48 well plate, and transfer each Gastruloid set to its appropriate secondary solution

7.3 Cover in aluminium and incubate 24:00:00 , 4 °C , on a orbital shaker

Mounting (D3)

8 Washes (to remove 2ry Ab):

8.1 Wash Gastruloids by transferring them to a well of a 6-well plate filled with 3mL PBS+FT

Note

Previous versions of this protocol call for PBS-/- instead of PBS-FT. There is a much higher increase of Gastruloids still sticking to the bottom of the wells in this case however. PBS-FT seems to be the safest option.

Safety information

You will need a separate PBS-FT well for each different sample condition, do not mix all Gastruloids into the same well

8.2 Wait for 00:20:00 min

8.3 Repeat for two more washes go to step #8.1 (total: 01:00:00 h)

Note

Again, if short on time, it is not necessary to wait 20 minutes for each wash. Gastruloids can be transferred serially across three PBSFT-filled wells, and left 20 minutes only in the last one.

9 Mounting:

- 9.1 Take a microscope slide, and add a drop of ~30uL of Fluoromount-G (mounting medium) to the centre of it

Note

While we usually do not, a spacer can be used to separate the sample from the coverslip, and to preserve the exact morphology of the sample. The absence of a spacer is not particularly destructive for early Gastruloids, but some damage can be seen for 168h and later timepoints.

The spacer can be either bought commercially, or made with two layers of double-sided sticky tape. The drop of mounting medium would then be added within the area defined by the spacer.

- 9.2 Transfer Gastruloids to the drop (see technique in "Guidelines")

Note

You can transfer any number of Gastruloids to the same drop. Depending on the size of their size, this might however increase the chance that they cluster together (e.g. 144h, 168h). If this happens, you can try to separate them and distribute them more evenly with an eyelash tool, being careful not to damage them.

When big Gastruloids are used, it is generally recommended to do multiple slides with fewer Gastruloids, rather than one with many all clumping together.

9.3 Gently drop a coverslip on top of the samples

Note

To avoid trapping air bubble, deposit a small drop of mounting medium at the centre of the coverslip prior to use.

9.4 Using the edge of a Kimtech wipe, absorb out excess liquid seeping out of the coverslip edges

9.5 Seal all sides of the coverslip with Permout hardening resin (we filled an empty nailpolish bottle with Permout, and use the brush for handling)

Note

Nail polish can be used instead of Permout, but there are reports of alcohol possibly being able to seep into the mounting medium and reducing the long-term shelf-life of the sample.

9.6 Keep the slide in a box away from light, 4 °C , until ready for imaging

NOTE: slides can be kept at 4 °C for months before imaging with little apparent loss of signal. Try however to image slides as close as possible to when they were made