VERSION 4

NOV 25, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bp3wmqpe

External link:
<https://doi.org/https://doi.org/10.1101/628974>

Collection Citation: Chen Chen, Lilan Hao, Weixia Wei, Fei Li, Liju Song, Xiaowei Zhang, Juanjuan Dai, Zhuye Jie, Jiandong Li, Xiaolei Song, Zirong Wang, Zhe Zhang, Liping Zeng, Hui Du, Huiru Tang, Tao Zhang, Huanming Yang, Jian Wang, Susanne Brix, Karsten Kristiansen, Xun Xu, Ruifang Wu, Huijue Jia 2020. Protocols for "The female urinary microbiota in relation to the reproductive tract microbiota." protocols.io <https://dx.doi.org/10.17504/protocols.io.bp3wmqpe> Version created by Lilan Hao

Protocols for "The female urinary microbiota in relation to the reproductive tract microbiota." V.4

Chen Chen¹, Lilan Hao¹, Weixia Wei², Fei Li¹, Liju Song¹, Xiaowei Zhang¹, Juanjuan Dai², Zhuye Jie¹, Jiandong Li¹, Xiaolei Song¹, Zirong Wang¹, Zhe Zhang¹, Liping Zeng², Hui Du², Huiru Tang², Tao Zhang¹, Huanming Yang¹, Jian Wang¹, Susanne Brix³, Karsten Kristiansen¹, Xun Xu¹, Ruifang Wu², Huijue Jia¹

¹BGI-Shenzhen, Shenzhen 518083, China.;

²Peking University Shenzhen Hospital, Shenzhen 518036, China.;

³Department of Biotechnology and Biomedicine, Technical University of Denmark, Soltofts Plads, Building 221, 2800 Kgs. Lyngby, Denmark.

GigaScience Press

BGI

Lilan Hao

ABSTRACT

Human urine is traditionally considered to be sterile, and whether the urine harbours distinct microbial communities has been a matter of debate. Potential links between female urine and reproductive tract microbial communities is currently not clear. Here, we collected urine samples from 147 Chinese women of reproductive age and explored the nature of colonization by 16S rRNA gene amplicon sequencing, quantitative real-time PCR, and live bacteria culture. To demonstrate the utility of this approach, the intra-individual Spearman's correlation was used to explore the relationship between urine and multiple sites of the reproductive tract. PERMANOVA was also performed to explore potential correlations between the lifestyle and various clinical factors and urinary bacterial communities. Our data demonstrated distinct bacterial communities in urine, indicative of a non-sterile environment. *Streptococcus*-dominated, *Lactobacillus*-dominated, and diverse type were the three most common urinary bacterial community types in the cohort. Detailed comparison of the urinary microbiota with multiple sites of the female reproductive tract microbiota demonstrated that the urinary microbiota were more similar to the microbiota in the cervix and uterine cavity than to those of the vagina in the same women. Our data demonstrate the potential connectivity among microbiota in the female urogenital system and provide insight and resources for exploring diseases of the urethra and genital tract.

MANUSCRIPT CITATION:

Chen C, Hao L, Wei W, Li F, Song L, Zhang X, Dai J, Jie Z, Li J, Song X, Wang Z, Zhang Z, Zeng L, Du H, Tang H, Zhang T, Yang H, Wang J, Brix S, Kristiansen K, Xu X, Wu R, Jia H, The female urinary microbiota in relation to the reproductive tract microbiota. GigaByte doi: <https://doi.org/10.1101/628974>

License: This is an open access collection distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Nov 25, 2020

Last Modified: Nov 25, 2020

COLLECTION integer ID:
44886

Protocol

NAME

DNA extraction for human microbe samples.

VERSION 1

CREATED BY

Lilan Hao

[OPEN](#) →

Protocol

NAME

A Bioinformatics Analysis workflow for 16S rRNA Amplicon Sequencing data

VERSION 1

CREATED BY

Lilan Hao

[OPEN](#) →

Protocol

NAME

Quantitative real-time PCR for the four Lactobacillus species

VERSION 1

CREATED BY

Lilan Hao

[OPEN](#) →