FEB 23, 2021

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.bspfdjn](https://doi.org/10.17504/protocols.io.bspfdjn)

External link:
<https://doi.org/https://doi.org/10.1101/2020.06.29.176958>

Collection Citation: Tianlin Pei, Mengxiao Yan, Yu Kong, Hang Fan, Jie Liu, Mengying Cui, Yumin Fang, Binjie Ge, Jun Yang, Qing Zhao 2021.

Protocols for "The genome analysis of *Tripterygium wilfordii* reveals TwCYP712K1 and TwCYP712K2 responsible for oxidation of friedelin in celastrol biosynthesis pathway". **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bspfdjn>

MANUSCRIPT CITATION:
 Pei T, Yan M, Kong Y, Fan H, Liu J, Cui M, Fang Y, Ge B, Yang J, Zhao Q, The genome of and characterization of the celastrol biosynthesis pathway. GigaByte doi: <https://doi.org/10.1101/2020.06.29.176958>

Protocols for "The genome analysis of *Tripterygium wilfordii* reveals TwCYP712K1 and TwCYP712K2 responsible for oxidation of friedelin in celastrol biosynthesis pathway"

GigaByte

Tianlin Pei^{1,2}, Mengxiao Yan¹, Yu Kong¹, Hang Fan^{1,2}, Jie Liu¹, Mengying Cui¹, Yumin Fang¹, Binjie Ge¹, Jun Yang^{1,2}, Qing Zhao^{1,2}

¹Shanghai Key Laboratory of Plant Functional Genomics and Resources, Shanghai Chenshan Botanical Garden, Shanghai Chenshan Plant Science Research Center, Chinese Academy of Sciences, Shanghai, China;

²State Key Laboratory of Plant Molecular Genetics, CAS Center for Excellence in Molecular Plant Sciences, Shanghai Institute of Plant Physiology and Ecology, Chinese Academy of Sciences, Shanghai, China

GigaScience Press

BGI

tianlinpei

ABSTRACT

Tripterygium wilfordii is a vine from the Celastraceae family that is used in traditional Chinese medicine (TCM). The active ingredient celastrol is a friedelane-type pentacyclic triterpenoid with putative roles as antitumor, immunosuppressive, and antiobesity agent. Here, we report a reference genome assembly of *T. wilfordii* with high-quality annotation by using a hybrid sequencing strategy, obtaining a 340.12 Mb total genome size, a contig N50 reaching 3.09 Mb. We successfully anchored 91.02% of the sequences into 23 pseudochromosomes using Hi-C technology and the superscaffold N50 reached 13.03 Mb. In addition, we annotated 31,593 structural genes, and a repeat percentage of 44.31%. Comparative evolutionary analyses showed that *T. wilfordii* diverged from species of Malpighiales approximately 102.4 million years ago. Based on integration of the genome, transcriptome and metabolite analyses, as well as *in vivo* and *in vitro* enzyme assays of the two CYP450 genes *TwCYP712K1* and *TwCYP712K2* the second biosynthesis step of celastrol was investigated. Syntenic analysis revealed that *TwCYP712K1* and *TwCYP712K2* were derived from a common ancestor. These results provide insights for further investigating pathways related to celastrol and valuable information to aid the conservation of resources. In addition, they helped us understand the evolution of Celastrales.

License: This is an open access collection distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 23, 2021

Last Modified: Feb 23, 2021

COLLECTION integer ID:
47559

Keywords: Tripterygium wilfordii, Genome, Celastrol, CYP450

Protocol

NAME

Irys NLRS DNA labeling and Data collection

VERSION 1

CREATED BY

GigaScience DatabaseGigaScience

[OPEN](#) →

Protocol

NAME

The pipeline of Hi-C assembly of the Scapharca broughtonii genome

VERSION 1

CREATED BY

Chang-Ming CM BaiYellow Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences

[OPEN](#) →

Protocol

NAME

Total DNA extraction from plant tissue using CTAB method

VERSION 1

CREATED BY

Robert Auber

[OPEN](#) →