APR 25, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.prydm7w

Protocol Citation: Kwame Agyemang Adusei, Alex K Owusu-Ofori 2018. PROTOCOLS FOR PREPARATION OF BLOOD SMEARS, REPORTING OF MALARIA PARASITES, PROCESSING OF BLOOD SAMPLES USING MALARIA RDT (HRP-2), BLOOD GROUPING AND DESIGN OF QUESTIONNAIRE. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.prydm7w>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Apr 25, 2018

Last Modified: Apr 25, 2018

PROTOCOLS FOR PREPARATION OF BLOOD SMEARS, REPORTING OF MALARIA PARASITES, PROCESSING OF BLOOD SAMPLES USING MALARIA RDT (HRP-2), BLOOD GROUPING AND DESIGN OF QUESTIONNAIRE

Kwame Agyemang Adusei¹, Alex K Owusu-Ofori²

¹Department of Medical Laboratory Service, School of Allied Health Sciences, Kwame Nkrumah University of Science and Technology ,Kumasi, Ghana .Tel: +233 501365180, Email: kwameagadusei@gmail.com;

²Department of Clinical Microbiology, School of Medical Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana. Tel: +233 209149370, Email: owusu_ofori@hotmail.com

Kwame Adusei

ABSTRACT

LABORATORY PROTOCOLS FOR PREPARATION OF BLOOD SMEARS, REPORTING OF MALARIA PARASITES, PROCESSING OF BLOOD SAMPLES USING MALARIA RDT (HRP-2), BLOOD GROUPING AND DESIGN OF QUESTIONNAIRE (PREVALENCE OF *PLASMODIUM* PARASITAEMIA IN BLOOD DONORS AND A SURVEY OF THE KNOWLEDGE AND PRACTICES OF TRANSFUSION MALARIA AMONG HEALTH WORKERS IN A HOSPITAL IN KUMASI, GHANA

MATERIALS

MATERIALS

Monoclonal antisera A,B and D **Contributed by users**

Keywords: Blood donors,
microscopy, rapid diagnostic
test (HRP-2), blood films,
blood groups

PREPARATION OF THICK SMEAR FOR MICROSCOPY

- 1 A drop of well mixed blood (6ul) was placed on a clean grease free slide using a pipette.
- 2 An applicator was used to spread the drop of blood in a circular way to ensure the spread of blood with moderate thickness, covering evenly an area about 1cm diameters.
- 3 Slides were labeled with subject's code number and allowed to air-dry.
- 4 The smears were stained with a freshly prepared 10% Giemsa stain for 15 minutes and washed with clean distilled water.
- 5 The slides were then placed on a drying rack in a vertical position and allowed to air dry.

PREPARATION OF THIN SMEAR FOR MICROSCOPY

- 6 A drop of well mixed blood (2ul) was placed about 1cm from the frosted end of a slide using a pipette.
- 7 A smooth clean edge of a second slide (spreader) was placed on the specimen slide, just in front

of the blood drop.

- 8 The spreader slide was held at 30° angle and was drawn back against the drop of blood.
- 9 As the blood started to spread to the ends of the spreader, the spreader was pushed forward with a light, smooth, and quick steady motion.
- 10 The frosted edge was labeled with the subject's code number and the blood smear was dipped in a Coplin jar containing absolute methanol for 10 seconds and then air dried again.
- 11 The smear was then stained with a freshly prepared Giemsa stain solution for 30 minutes and washed with distilled water
- 12 The slides were then placed on a drying rack and allowed to air dry. An oil immersion was put on the prepared thick film and was examined using the X100 objective lens and the thin films were first examined using the X40 with an oil immersion placed to the feathery tip of the smear to check the staining, morphology and distribution of the cells and afterwards the objective lens is changed to X100 to examine the malaria parasites.

REPORTING OF MALARIA PARASITES

- 13 Number of parasites seen were counted in relation to a predetermined number of white blood cells and an average of 8000/ μ L was taken as standard. 200 white blood cells were counted in 100 fields. All parasite species and forms including both sexual and asexual forms were counted together.
- 14 $\text{NUMBER OF PARASITES}/\mu\text{L} = \text{Number of parasites counted} / \text{Number of white blood cells counted} \times 8000$

BLOOD SAMPLES PROCESSING FOR MALARIA DIAGNOSIS U...

- 15 In the laboratory, 5ul of whole blood from the labeled EDTA tube, was taken with a pipette provided with the test kit.
- 16 The blood was pipetted into the round sample well of the RDT.
- 17 Four drops of assay diluent were placed vertically into the square assay diluent well and the results were read after 30 minutes.
- 18 The same procedure was performed on samples collected from blood donors and non-donors.
- 19 A positive results was recorded when the test and control bands appeared after 30 minutes. A negative results was also recorded when only the control band showed.

BLOOD GROUPING

- 20 The tile method using the forward grouping was used for determining the blood groups of blood donors and non-donors (controls).
- 21 5ul of blood donors and non-donors blood samples were put in the appropriate square of the tile.
- 22 A drop of a known antisera was added to the sample.

- 23** An applicator was used in mixing and the tile was rocked forward and backward for 3 minutes. The results were read macroscopically upon agglutination and in cases where the results were unclear, were read microscopically

QUESTIONNAIRE

- 24** The questionnaire was categorized into a three-part document with a socio-demographics section, knowledge section and a section on attitudes and practices. Responses to a total of 20 opened and closed complete questions were unspecified so as to obtain validated answers without fear of victimization. Health workers were given less than 24hours to fill the questionnaire and those who did not submit were followed up in person