OCT 02, 2020

Purification of Haustoria from Arabidopsis in Response to Infection by *E. cichoracearum*

Harley King¹

¹LuminUltra

Harley King Workspace

Harley King
LuminUltra

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.tqkemuw

Protocol Citation: Harley King 2020. Purification of Haustoria from Arabidopsis in Response to Infection by *E. cichoracearum*. [protocols.io](https://dx.doi.org/10.17504/protocols.io.tqkemuw) <https://dx.doi.org/10.17504/protocols.io.tqkemuw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
We are still developing and optimizing this protocol

Created: Sep 18, 2018

Last Modified: Oct 02, 2020

PROTOCOL integer ID:
15852

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

This procedure describes steps for the purification of *Arabidopsis thaliana* haustoria in response to powdery mildew infection by fungus *Golovinomyces cichoracearum*.

Grind Plants Containing Haustoria

1. Weigh out 5-10g frozen Arabidopsis leaves. Set aside 2-3 leaves for quantifying haustoria with

- a confocal microscope before purification.
2. Grind leaves in ice-cold 8 mL 1x PBS buffer pH 7.4 for 1 min in kitchen blender in cold room.
 3. Filter grindate through 100um Nylon mesh. Haustoria are about 10-20um in diameter.
 4. With a glass rod, remove unfiltered debris to blender. PBS buffer can be used to wash the membrane with a transfer pipette into the blender.
 5. Add 5 mL 1x PBS buffer and grind for another 45 seconds.
 6. Filter through 100um Nylon mesh.
 7. Filter the collected filtrate from steps 3 and 6 and pass through a 40um mesh. A vacuum and modified filter assembly can be used to expedite the purification. Wash the membrane with PBS buffer into the trash.
 8. Pass filtrate through 40um mesh again.
 9. Transfer ~13-15mL to a 15mL conical tube. Pellet filtrate at 1000g for 5 min.
 10. Remove supernatant.
 11. Resuspend pellet in 2mL 1x PBS buffer. Set aside 40ul for quantifying haustoria.
 12. Investigate haustoria with confocal microscope and a hemacytometer before committing to the next steps.

Note

Col-O plants should be processed concurrently as a negative control especially if haustoria will be filtered by FACS.

Purify Haustoria Using Percoll Cushion

- 2 Using Percoll, make a 60% solution and a 40% solution with 1x PBS.
Layer 4mL of the 60% solution on the bottom of a 15mL conical tube.
Layer 6mL of the 40% solution on the top of the 60% solution.
Gently add 1mL of resuspended pellet containing the haustoria to the top 40% Percoll solution.
Centrifuge at 2500g for 10 minutes.
The haustoria will penetrate the 40% layer but not the 60% layer.
Remove the 40% layer. Mix, then pellet at 1000g for 5 min.
Resuspend in 2mL. Remove 40ul aliquot for quantifying haustoria.
If sufficient, proceed to next step.

Incubate Haustoria with anti-GFP (Chromotek) Conjugated ...

- 3
 1. Conjugate anti-GFP Trap antibody to Dynabeads following manufacturer's recommended protocol.
 2. Incubate beads with haustoria with end-over-end rotation overnight.
 3. Wash beads 5x with 1x PBS pH7.4.

-
4. Disrupt haustoria-GFP Trap binding following manufacturer's protocol.
 5. Proceed to FACS purification

4 Follow previous experiments to use as a baseline in purifying haustoria with FACS. Use col-o plants as negative control.