APR 14, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.u7ueznw

External link:
<http://www.bcell.org>

Protocol Citation: Nancy Smith, Haopeng Lin, Mourad Ferdaoussi, Patrick E Macdonald 2020. Purification of Mouse Pancreatic Islets using Histopaque Gradient Centrifugation. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.u7ueznw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 31, 2018

Last Modified: Apr 14, 2020

PROTOCOL integer ID:
 17364

Purification of Mouse Pancreatic Islets using Histopaque Gradient Centrifugation

Nancy Smith¹, Haopeng Lin¹, Mourad Ferdaoussi¹,
 Patrick E Macdonald¹

¹University of Alberta

CIRTNR2FIC

Nancy Smith
 University of Alberta

ABSTRACT

This protocol details the purification of mouse pancreatic islets using histopaque gradient centrifugation.

MATERIALS

MATERIALS

- ☒ Hank Balanced Salt Solution 1X **Sigma Aldrich Catalog #H6136**
- ☒ Albumin Bovine Serum (30%) **equitech bio, inc. Catalog #160928-0262**
- ☒ Histopaque 1077 **Sigma Aldrich Catalog #1077-1**
- ☒ RPMI 1640 (11.1mM glucose) **Gibco - Thermo Fischer Catalog #11875-119**
- ☒ FBS Canadian Origin **Gibco - Thermo Fischer Catalog #12483-020**
- ☒ Cell dissociation sieve **Sigma Aldrich Catalog #CD1-1KT**
- ☒ 50ml Falcon tubes **Corning Catalog #352070**
- ☒ 30ml syringe **BD Biosciences Catalog #302832**
- ☒ 18G needle **BD Biosciences Catalog #305196**
- ☒ Penicillin Streptomycin **Invitrogen - Thermo Fisher Catalog #15140 122**
- ☒ Sodium bicarbonate **Sigma Aldrich Catalog #S5761**

BEFORE START INSTRUCTIONS

1X Hanks' Balanced Salt Solution (HBSS) + BSA, Coating and Mouse Islet Culture Media should be prepared in advance and should be cold before starting gradient steps.

Keywords: Histopaque,
islets, isolation, gradient,
mouse, pancreas

Solution Prep - 1X Hanks' Balanced Salt Solution (HBSS) + ...

- 1 Measure out 900ml of room temperature H₂O
900 mL MiliQ water
- 2 While gently stirring the water, add the powdered medium. Stir until dissolved. DO NOT HEAT.
- 3 Rinse original vial with water to remove traces and add to above.
- 4 Add 0.35g sodium bicarbonate and stir until dissolved.
0.35 g Sodium Bicarbonate
- 5 Adjust pH to 7.4
- 6 Bring solution to 1L
- 7 Take 34ml of the prepared 1X HBSS and add 6ml of 30% BSA (equitech bio, inc 160928-0262) in a 50ml falcon tube. This is to be used for coating.
- 8 Add 4ml of 30% BSA (equitech bio, inc 160928-0262) to the prepared 1X HBSS

9 Store at 4°C

Solution Prep - Mouse Islet Culture Media

10 Add FBS and Pen/Strep to bottle of RPMI

500ml RPMI 1640 (11.1mM glucose)	Gibco 11875-119
50ml FBS Canadian Origin	Gibco 12483-020
5ml Pen/strep (10000 Unit/ml/10000 ug/ml)	Gibco 15140-122

11 Store at 4°C

Gradient Protocol

12 Once digest of pancreas in the waterbath is complete (see Mouse Pancreatic Islet Isolation protocol, Step 16)

<https://www.protocols.io/view/mouse-pancreatic-islet-isolation-sqaedse>

Shake 50ml falcon tube 30 times.

13 Bring up volume to 50ml by adding 1X HBSS+BSA to falcon tube.

14 Centrifuge for 2 minutes at 339 g at 4°C

- 15** Remove supernatant
- 16** Re-suspend pellet in 20ml of 1X HBSS+BSA. Pour re-suspended pellet through #60 screen (cell dissociation sieve, Sigma CD1-1KT) that has been rinsed with coating. Using a 30ml syringe filled with 1X HBSS+BSA and 18G needle rinse screen (use about 20ml). Return islets to 50ml tube and rinse the beaker with the remaining 10ml of 1X HBSS+BSA, add it to the tube.
- 17** Centrifuge for 2 minutes at 339 g at 4°C
02:00:00
4 °C
- 18** Remove supernatant. Add 10ml 1X HBSS+BSA, re-suspend pellet. Bring up the volume to 50ml with 1X HBSS+BSA.
- 19** Centrifuge for 2 minutes at 339 g at 4°C
02:00:00
4 °C
- 20** Remove supernatant. Add 10ml 1X HBSS+BSA, re-suspend pellet. Bring up the volume to 50ml with 1X HBSS+BSA.
- 21** Centrifuge for 2 minutes at 339 g at 4°C
02:00:00
4 °C
- 22** Remove supernatant. Leave tube upside down on papertowel to drip dry (30 seconds). Wipe the interior of the tube with a Kimwipe using a long forceps (do not touch the pellet). Leave tubes upright on ice.

23 Re-suspend the pellet with 5ml of Histopaque 1077 (Sigma 1077-1 500ml) with pipette that has been rinsed with coating. Add another 5ml of Histopaque and mix using coated pipette.

24 Lay the tube at an angle on ice. Add gradually 10ml of the Mouse Media to ensure that the two mediums do not mix.

25 Centrifuge for 30 minutes at 339 g at 4°C with the **brake turned off**.

30:00:00

4 °C

Note

Some centrifuges have an acceleration/deceleration speed setting. Use slowest rate of speed for both acceleration and deceleration .

26 Transfer the supernatant (both layers) into a new single 50ml falcon tube. Bring up to volume with 1X HBSS+BSA.

Note

Keep tubes with the pellet on ice. If islets are not in the the supernatant, they may be in the pellet and can be recovered by hand picking in Mouse Islet Media.

27 Centrifuge for 3 minutes at 339 g at 4°C.

03:00:00

4 °C

28 Decant the supernatant. Re-suspend the pellet in 10 ml of Mouse Media.

- 29 Hand pick islets if desired or necessary. Incubate islets in Mouse Islet Media at 37°C with 5% CO₂ until ready for use in experiment.