JUL 29, 2017

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.if3cbqn](https://doi.org/10.17504/protocols.io.if3cbqn)

External link:
<https://doi.org/10.1371/journal.pone.0181703>

Protocol Citation: Zhu Feng 2017. Purification of *Nosema bombycis* spores. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.if3cbqn>

MANUSCRIPT CITATION:
 Chen G, Wang W, Chen H, Dai W, Peng X, Li X, Tang X, Xu L, Shen Z (2017) Functional characterization of an aquaporin from a microsporidium, *Nosema bombycis*. PLoS ONE 12(7): e0181703. doi: [10.1371/journal.pone.0181703](https://doi.org/10.1371/journal.pone.0181703)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Purification of *Nosema bombycis* spores

PLOS One

Zhu Feng¹

¹Jiangsu University of Science and Technology

Chen Gong

ABSTRACT

Infected beetles were collected and homogenized in sterile water. The homogenates were filtered through four layers of cheesecloth and centrifuged at 3000g for 15 min. The pellets were resuspended in sterile water, and the spores were purified by Percoll gradient centrifugation using 90% Percoll at 15,000g for 40 min. The spore band was collected and washed several times with sterile water. The purified spores (n = 50) were measured under a light microscope (IL/Leica Microsystems, Inc., Deerfield) with an ocular micrometer and photographed with the Microscope USB Camera.

ATTACHMENTS

[A new isolate of *Nosema* sp. \(Microsporidia, Nosematidae\) from.pdf](#)

Created: Jun 15, 2017

Last Modified: Mar 19, 2018

PROTOCOL integer ID:
6363