AUG 13, 2020

DOI:
dx.doi.org/10.17504/protocols.io.bjnsknee

Protocol Citation: Angel A Justiz-Vaillant 2020. Purification of the any avian IgY using chloroform, trichloroacetic acid and affinity purification using HiTrap™ Columns.. [protocols.io https://dx.doi.org/10.17504/protocols.io.bjnsknee](https://dx.doi.org/10.17504/protocols.io.bjnsknee)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 13, 2020

Last Modified: Aug 13, 2020

PROTOCOL integer ID:
 40370

🌐 Purification of the any avian IgY using chloroform, trichloroacetic acid and affinity purification using HiTrap™ Columns.

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to protocols.io is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with protocols.io, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

This IgY purification technology can be used to purified IgY from the egg yolk of any bird as ducks, bantam hens, ostrich, chicken, pigeon, pheasant, quail, guinea hen, cattle egret and goose.

MATERIALS

MATERIALS

⊗ 500ml Trichloroacetic acid (TCA) Solution, 10% (w/v) **G-Biosciences Catalog #786-886**

⊗ Chloroform **Sigma Catalog #319988**

⊗ PBS **Life Technologies**

⊗ Affinity Purification Using HiTrap™ Columns **Contributed by users**

- 1 The IgY fraction of bantam hen was isolated by the chloroform-Trichloroacetic acid (TCA) -affinity chromatography method.
- 2 The egg is washed with warm water and the egg yolk is separated from the egg white.
- 3 The membrane is broken and the egg yolk collected.
- 4 The egg yolk is then diluted 1:3 in phosphate buffered saline (PBS), pH 7.4.
- 5 To 1/3 of the egg yolk mixture an equal volume of chloroform is added,
- 6 The mixture is then shaken and centrifuged for 30 min (1000×g, RT).
- 7 The supernatant is decanted and drop 1 ml of cold TCA and let the sample stand for 30 min at RT.

- 8 The mixture is then centrifuged as previously described.
- 9 The precipitate containing IgY is dissolved in PBS (pH 7.4) at a volume equivalent to 1/6 of the original volume of the egg yolk.
- 10 After that the preparation is dialyzed against 1L of PBS (pH: 7.4 for 24 h at 4°C).
- 11 The IgY is then removed from the dialysis tubing.
- 12 IgY concentration is assessed by an available methodology.
- 13 IgY samples then are further purified by affinity chromatography.
- 14 The preparation can be stored at –20°C.
- 15 By a sandwich ELISA the IgY concentration can be assessed.

- 16** Use Affinity Purification Using HiTrap™ Columns as follows:
- 17** Fill the syringe with distilled water.
- 18** Connect the column to the syringe
- 19** Snap off the tab on the column outlet.
- 20** Wash out the ethanol with distilled water.
- 21** Equilibrate the column with binding buffer.
- 22** The recommended flow rate is 5 ml/min*.
- 23** Apply the sample using a syringe linked or fitted to Luer connector
- 24** Use a flow rate of 1 to 5 ml/min during sample application

- 25** Wash with at least 51 ml of binding buffer.
- 26** Maintain a flow rate of 10 ml/min for washing.
- 27** Elute with 50 ml of elution buffer.
- 28** After elution, regenerate the column by using wash buffer
- 29** Re-equilibrate the column with 26 ml of binding buffer.
- 30** Store it until a new purification