



Jul 19, 2024

qPCR based multipathogen detection for Salmonella Paratyphi "A" and Vibrio cholerae from wastewater samples

DOI

<https://dx.doi.org/10.17504/protocols.io.6qpvr8m8plmk/v1>

Dilip Abraham¹, Blossom Benny Sam¹, Nirmal Kumar¹, Raju Ravi¹, Venkata Raghava Mohan²

¹Wellcome Trust Research Laboratory, Christian Medical College Vellore, India;

²Department of Community Health and Development, Christian Medical College Vellore, India



Dilip Abraham

CMC Vellore

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.6qpvr8m8plmk/v1>

Protocol Citation: Dilip Abraham, Blossom Benny Sam, Nirmal Kumar, Raju Ravi, Venkata Raghava Mohan 2024. qPCR based multipathogen detection for Salmonella Paratyphi "A" and Vibrio cholerae from wastewater samples. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.6qpvr8m8plmk/v1>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited



Protocol status: Working

We use this protocol and it's working

Created: July 11, 2024

Last Modified: July 19, 2024

Protocol Integer ID: 103282

Keywords: qPCR, Environmental surveillance, Wastewater samples, Salmonella Paratyphi A, Vibrio cholerae, multipathogen detection for salmonella paratyphi, different multiplex pcr assay, using multiplex qpcr assay, multiplex qpcr assays from tna, qpcr assay, vibrio cholerae from wastewater sample, vibrio cholerae gene target, salmonella paratyphi, time pcr assay, vibrio cholerae, validation of the primer probe combination, qpcr, primer probe combination, pcr, based multipathogen detection, wastewater sample

Funders Acknowledgements:

Bill & Melinda Gates Foundation

Grant ID: INV-049093

Bill & Melinda Gates Foundation

Grant ID: INV-048134

Abstract

The following protocol is optimized for the detection of *Salmonella* Paratyphi A and *Vibrio cholerae* using multiplex qPCR assays from TNA extracted from environmental samples (wastewater samples). Total Nucleic Acid (TNA) extracted from the pellet particles is used as template in a real-time PCR assay which uses primer-probes specific for *Salmonella* Paratyphi A or *Vibrio cholerae* gene targets. Three different multiplex PCR assays were designed following standardization and validation of the primer probe combinations.



Guidelines

This protocol describes qPCR based detection of *Salmonella* Paratyphi "A" and *Vibrio cholerae* gene targets in three PCR panels from TNA extraction of processed environmental samples that were spiked with internal controls (SPC EGT control DNA - referenced below) . Additionally, HF183 which serves as a marker of fecal contamination in environmental samples, is included in the qPCR assays. Due to the similarity in cycling conditions, these assays can be conducted simultaneously for a sample as three distinct panels.

The positive controls (PCs) used here are linear oligos (gBlocks) that have been used in generating standard curves. Ideally the concentration of PCs used should yield Ct values that fall in the linear phase of the amplification curve.

Protocol

NAME

Extraction of Total Nucleic Acid from Environmental Samples for the Detection of Bacterial and Viral Targets

CREATED BY

Dilip Abraham

Preview



Materials

1. QuantStudio7 Flex machine
2.  Platinum[®] Quantitative PCR SuperMix-UDG **Thermo Fisher Catalog #11730025**
3.  qPCR DNA Extraction and Inhibition Control CY5-QXL670 **Eurogentec Catalog #RT-SPCC-Q02**
4.  Nuclease-free water **Ambion Catalog #AM9932**
5. Primers and Probes (Sigma/IDT)(detailed in protocol)
6. Positive Controls: gBlocks gene fragments (IDT) (detailed in protocol)
7. MicroAmp[™] Optical 96-Well Reaction Plate with Barcode Thermofisher Catalog #4306737
8. MicroAmp[™] Optical Adhesive Film Thermofisher Catalog # 4311971
9. 1.7 mL MaxyClear Snaplock Microcentrifuge Tube Axygen Catalog #MCT-175-C
10. Finnpiquette F1 100 to 1000 µL Thermo Fisher Catalog #4641100N
11. Finnpiquette F1 20 to 200 µL Thermo Fisher Catalog #4641080N
12. Finnpiquette F1 2 to 20 µL Thermo Fisher Catalog #4641060N
13. Finnpiquette F1 0.2 to 2 µL Thermo Fisher Catalog #4641010N
14. ART Barrier Specialty Pipette tips 1000 µL Thermo Fisher Catalog #2279-05PK
15. ART Barrier Specialty Pipette tips 200 µL Thermo Fisher Catalog #2069-05PK
16. ART Barrier Specialty Pipette tips 20 µL Thermo Fisher Catalog #2149P-05PK
17. ART Barrier Specialty Pipette tips 10 µL Thermo Fisher Catalog #2139-05PK
18. Microplate Centrifuge, PCR Plate Spinner VWR[®] Catalog #VWRU89184-610

Before start

Ensure that the fluorescent dye used for TaqMan probes is compatible with the machine being used and properly calibrated for it.

Primer-Probe panel

- 1 The following primers and probes are employed for the detect *Salmonella* Paratyphi A and *Vibrio cholerae* specific gene targets.

1.1 PCR 1: *Salmonella* Paratyphi A (SPA2308) and *V. cholerae* ctxA PCR panel

A	B	C
TARGET	PRIMER S	SEQUENCES 5' TO 3'
ctxA	Forward	TCCGGAGCATAGAGCTTGGA
	Reverse	TCGATGATCTTGGAGCATTCC
	Probe	[CY5] - CCGTGGATTCATCATGCACCGC - [BHQ2]
SPA2308	Forward	ACGATGATGACTGATTTATCGAAC
	Reverse	TGAAAAGATATCTCTCAGAGCTGG
	Probe	[FAM] - CCCATACAATTTTCATTCTTATTGAGAATGCGC - [BHQ1]

Table1: SPA and ctxA Primer and probe sequences. Fluorescent dyes and quenchers are shown in square brackets.

1.2 PCR 2: *Vibrio cholerae*_1 PCR panel

A	B	C
TARGET	PRIMERS	SEQUENCES 5' TO 3'
HF183	Forward	ATCATGAGTTCACA GTCCG
	Reverse	CTTCCTCTCAGAACCCCTATCC
	Probe	[FAM] - CTAATGGAACGCATCCC - [BHQ1]
wbfO139	Forward	AGAAGCCAGTCGCAGTAAAG
	Reverse	TCGCCATCTTCCAGCATAAA
	Probe	[TAMRA] - TGGTGGTACAGCTTAGCCGCATTA - [BHQ2]

	A	B	C
	tcpA classic	Forward	GCGTAATGCAGCAGCTAATAAA
		Reverse	TATGGGAACATATCACCGACAC
		Probe	[JOE] - ATGGTCTGACACAGGCTCAATGCA - [BHQ1]

Table2: *Vibrio cholerae*_1 PCR Primer and probe sequences. Fluorescent dyes and quenchers are shown in square brackets.

1.3 PCR 3: *Vibrio cholerae*_2 PCR panel

	A	B	C
	TARGET	PRIMER S	SEQUENCES 5' TO 3'
	OmpW	Forward	TCAATGATAGCTGGTTCCTCAAC
		Reverse	CGATGATAAATACCCAAGCATTGA
		Probe	[JOE]_TGGTATGCCAATATTGAAACAACG_[BHQ1]
	wbeO1	Forward	GTTGAGAAGGGCGGTCTAATAA
		Reverse	TGTCTGGTACTTGAGTTGGTAAG
		Probe	[FAM] - TGCCTCAGCAATGGA - [BHQ1]
	tcpA Eltor	Forward	ATCCTTTCACTGGTACAGCTATG
		Reverse	GTCAAGCCACCGACTGTAAT
		Probe	[TAMRA]- ACGAACTCTGCAGCGAATAAAGC-[BHQ2]

Table3: *Vibrio cholerae*_2 PCR Primer and probe sequences. Fluorescent dyes and quenchers are shown in square brackets.

Primer-probe reconstitution

- 2 To reconstitute the lyophilized primers/probes, use the nmole information on the specification sheet received with primers/probes.
- 2.1 Multiply nmol value by 10 to get the required volume of Nuclease Free Water (NFW) needed to reconstitute the lyophilized primer/probes.

Ex: For a primer with 30 nmoles, to make 100 micromolar (μM) stock solution:
 $30 \text{ nmol} \times 10 = 300 \mu\text{L}$ of NFW to make 100 micromolar (μM) stock solution.

2.2 Add the required volume of nuclease free water, pulse vortex and spin down. This is the primer/probe stock with 100 micromolar (μM) concentration.

2.3 Store at -20°C for long term storage.

Primer-probe dilution

3 Prepare a working stock from 100 micromolar (μM) stock solution.

3.1 In a fresh tube add 10 μL of 100 micromolar (μM) primer/probe stock and 90 μL of nuclease free water to give 100 μL of 10 micromolar (μM) working primer/probe.

3.2 Store at 4°C for frequent usage or -20°C for long-term storage.

qPCR controls

4 **Positive control:** gBlocks gene fragments corresponding to each gene target is included in qPCR assay to use as positive control.

A	B	C	D
gBlock gene	Sequence(5'-3')	Accession No:	bp size
SPA2308	ACGATGATGACTGATTTATCGAACAACGAC TCTCCCATACAATTTTCATTCTTATTGAGAAT GCGCTTATGTAATTTATACCCAGCTCTGA GAGATATCTTTTCA	FM200053.1	105
ctxA	TCCGGAGCATAGAGCTTGGAGGGAAGAGC CGTGGATTCATCATGCACCGCCGGGTTGT GGGAATGCTCCAAGATCATCGA	AF463401.1	80
HF183	GGGATCATGAGTTCACATGTCCGCATGATT AAAGGTATTTTCCGGTAGACGAT GGGGATGCGTTCCATTAGATAGTAGGCGG	MT464394.1	132

A	B	C	D
	GGTAACGGCCACCTAGTCAACG ATGGATAGGGGTTCTGAGAGGAAGGTC		
wbfO139	AGAAGCCAGTCGCAGTAAAGCACTAGGGC GCATGGTGGTACAGCTTAGCCGCATTATGC GAGATGAGCCGGGTGCGGATTTTATGCTG GAAGATGGCGA	AB012956.1	99
tcpA classic	GCGTAATGCAGCAGCTAATAAAGCATTTGC AATTTCAGTGGATGGTCTGACACAGGCTC AATGCAAGACACTTATTACCAAGTGTGCGGTG ATATGTTCCATA	M33514.1	102
OmpW	TCAATGATAGCTGGTTCCTCAACGCTTCTG TGTGGTATGCCAATATTGAAACAACGGCAA CCTACAAAGCAGGTGCAGATGCCAAATCC ACGGATGTTGAAATCAATCCTTGGGTATTT ATGATCG	X51948 modified	126
wbeO1	GTTGAGAAGGGCGGTCTAATAACACCTAA AGAGTTTGCAGAGAAGCTTGCCTCAGCAA TGGATAAGGCTCTTGTACGCTTACCAACTC AAGTACCAGACA	KC152957.1	100
tcpA El-Tor	ATCCTTTCACTGGTACAGCTATGGGGATTT TCTCATTTCCACGAACTCTGCAGCGAAT AAAGCATTGCAATTACAGTCGGTGGCTT GAC	KP187623.1	91

Table4: Sequences used for gBlocks gene fragments

- 5 **Negative control:**  3 µL of extraction blank of each batch of extraction.
NTC: Master mix alone used for no template control.

Quantitative PCR

- 6 Thaw qPCR reagents and samples on ice and briefly spin it down.
Prepare master mix for each of the 3 primer-probe panels.
- 7 **[PCR 1] *Salmonella* Paratyphi A (SPA2308) and ctxA**
- 7.1 Prepare the master mix as follows for the number of samples, positive and negative controls, NTC and one extra reaction to account for any pipetting error.



A	B
REAGENT	VOLUME FOR 1 REACTION (μL)
UDG Mix	12.5
MgCl ₂	1
Rox dye	0.05
ctxA F Primer (10 μM)	0.5
ctxA R Primer (10 μM)	0.5
ctxA Probe (10 μM)	0.25
SPA2308 F (10 μM)	0.5
SPA2308 R (10 μM)	0.5
SPA2308 Probe (10 μM)	0.25
NFW	5.95

Table5: Mastermix composition for SPA2308 and ctxA gene targets

8 [PCR 2] *Vibrio cholerae*_1 PCR

- 8.1 Prepare the master mix as follows for the number of samples, positive and negative controls, NTC and one extra reaction to account for any pipetting error.

A	B
REAGENT	VOLUME FOR 1 REACTION (μL)
UDG Mix	12.5
MgCl ₂	1
ROX dye	0.05
HF183 F Primer (10 μM)	0.5
HF183 R Primer (10 μM)	0.5

A	B
HF183 Probe (10 µM)	0.25
wbfO139 F Primer (10 µM)	0.5
wbfO139 R Primer (10 µM)	0.5
wbfO139 Probe (10 µM)	0.25
tcpA classic F Primer (10 µM)	0.5
tcpA classic R Primer (10 µM)	0.5
tcpA classic Probe (10 µM)	0.25
SPC (10 X EGT Control Mix)	2.5
NFW	2.2

Table6: Mastermix composition for *Vibrio cholerae*_1 PCR gene targets

Note

SPC is Sample Processing Control, an optimized TaqMan control designed to be used as qPCR DNA extraction and inhibition control. It is detected by Cy5-labelled probe (Cy5-QXL670 Probe) which comes as ready to use primer probe mix.

9 [PCR 3] *Vibrio cholerae*_2 PCR

- 9.1 Prepare the master mix as follows for the number of samples, positive and negative controls, NTC and one extra reaction to account for any pipetting error.

A	B
REAGENT	VOLUME FOR 1 REACTION (µL)
UDG Mix	12.5
MgCl ₂	1
ROX dye	0.05

A	B
OmpW F Primer (10 µM)	0.5
OmpW R Primer (10 µM)	0.5
OmpW Probe (10 µM)	0.25
wbeO1 F Primer (10 µM)	0.5
wbeO1 R Primer (10 µM)	0.5
wbeO1 Probe (10 µM)	0.25
tcpA El-Tor F Primer (10 µM)	0.5
tcpA El-Tor R Primer(10µM)	0.5
tcpA El-Tor Probe (10 µM)	0.25
NFW	4.7

Table7: Mastermix composition for *Vibrio cholerae*_2 PCR gene targets

- 10 For each PCR, aliquot 22 µl of master mix to each required well in a 96-well plate. Add 3µl of sample, or 3 µl of nuclease free water for negative controls.
- 11 Seal the plate with a roller sealer and then centrifuge the plate for 1 min at 2,000g.
- 12 Load the plate into Quantstudio7 flex instrument after proper initiation of the instrument. Open QS7 software, then select "New Experiment set up".
- 13 Set up the experiment properties with 96-well block, TaqMan reagents, 0.2 ml PCR plate and standard run. Define sample ID and define the targets as described for respective PCR panels. Assign targets and sample ID to each well.
- 14 Set up the PCR cycling method as described below. Cycling conditions remain the same for all 3 qPCRs.

A	B	C
Step (Hold Stage)	Temperature	Time
Reverse Transcription	50°C	2 MIN

A	B	C
PCR initial heat activation	95°C	2 MIN
2-step cycling (40 cycles)		
Denaturation	95°C	15 SEC
Combined annealing/extension	60°C (data collection step)	30 SEC

Start the qPCR by clicking “Run.”

- 15 Once the run is complete, adjust the thresholds and baseline if any abnormal baseline at the start or at the end is observed, which may lead to a false-positive curve. Verify if the PC is within the range using the cut-off Ct values chosen from running the standards.
- 16 Export the result to excel/csv file and upload both run and csv files.
- 17 The threshold for each target can be set such that the PC for that target falls within the pre-defined range obtained with the standard curves.

The sample is considered positive if the amplification curve is appropriate and the Ct value falls below the defined cut-off thresholds for each target.

A separate protocol, provided in the Typhoid ES workspace, serves as an example and can be followed to generate Ct cut-off values:

Protocol

NAME

Generating Ct cut-off values using gBlocks gene fragments

CREATED BY

Jonathan Rigby

Preview

Protocol references

- 1] Kahler AM, Haley BJ, Chen A, Mull BJ, Tarr CL, Turnsek M, Katz LS, Humphrys MS, Derado G, Freeman N, Boncy J, Colwell RR, Huq A, Hill VR. Environmental surveillance for toxigenic *Vibrio cholerae* in surface waters of Haiti. *Am J Trop Med Hyg.* 2015 Jan;92(1):118-25. doi: 10.4269/ajtmh.13-0601. Epub 2014 Nov 10. PMID: 25385860; PMCID: PMC4347365. doi: [10.4269/ajtmh.13-0601](https://doi.org/10.4269/ajtmh.13-0601)

- 2] Alejandro Garrido-Maestu, María-José Chapela, Elvira Peñaranda, Juan M. Vieites, Ana G. Cabado, In-house validation of novel multiplex real-time PCR gene combination for the simultaneous detection of the main human pathogenic vibrios (*Vibrio cholerae*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*), *Food Control*, Volume 37, 2014, Pages 371-379, ISSN 0956-7135, <https://doi.org/10.1016/j.foodcont.2013.09.026>.

- 3] David R. Greig, Tadgh J. Hickey, Megan D. Boxall, Hafsa Begum, Amy Gentle, Claire Jenkins, Marie A. Chattaway, A real-time multiplex PCR for the identification and typing of *Vibrio cholerae*, *Diagnostic Microbiology and Infectious Disease*, Volume 90, Issue 3, 2018, Pages 171-176, ISSN 0732-8893, <https://doi.org/10.1016/j.diagmicrobio.2017.11.017>.

- 4] Nga TV, Karkey A, Dongol S, et al. The sensitivity of real-time PCR amplification targeting invasive *Salmonella* serovars in biological specimens. *BMC Infect Dis.* 2010;10:125. Published 2010 May 21. doi:10.1186/1471-2334-10-125.

- 5] <https://www.eurogentec.com/assets/b2503b94-10d4-4572-b1fd-95969baeccc3/tds-en-rt-spcc-q-qpcr-dna-extraction-and-inhibition-control-cy5-qxl670.pdf>

- 6] Thong KL, Tham KBL, Ngoi ST, et al. Molecular characterization of *Vibrio cholerae* O1 El Tor strains in Malaysia revealed genetically diverse variant lineages. *Transbound Emerg Dis.* 2022;69(4):e693-e703. doi:10.1111/tbed.14368