VERSION 1

SEP 08, 2020

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bjrkkm4w

Protocol Citation: Adrieli Sachett¹, Matheus Gallas-Lopes¹, Greicy M M Conterato², Radharani Benvenuti¹, Ana P Herrmann¹, Angelo Piato 2020. Quantification of nonprotein sulfhydryl groups (NPSH) optimized for zebrafish brain tissue. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjrkkm4w>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Aug 14, 2020

Last Modified: Sep 08, 2020

PROTOCOL integer ID:
40460

Quantification of nonprotein sulfhydryl groups (NPSH) optimized for zebrafish brain tissue V.1

Adrieli Sachett¹, Matheus Gallas-Lopes¹,
Greicy M M Conterato², Radharani Benvenuti¹,
Ana P Herrmann¹, Angelo Piato¹

¹Universidade Federal do Rio Grande do Sul;

²Universidade Federal de Santa Catarina

Fish behavior and physiology

LAPCOM

Angelo Piato

Universidade Federal do Rio Grande do Sul

ABSTRACT

Zebrafish are increasingly used as a model animal in neuroscience research. Here we describe a protocol to quantify nonprotein sulfhydryl groups (NPSH), an indirect evaluation of the levels of reduced glutathione (GSH), a major oxidative stress defense in the central nervous system.

GUIDELINES

This protocol is intended to standardize nonprotein sulfhydryl groups quantification of zebrafish brain tissue samples. It can be adapted for other fish species.

Keywords: NPSH, Oxidative stress, Zebrafish brain tissue

MATERIALS

MATERIALS

☒ Gloves **Contributed by users**

☒ 96 well plate **Contributed by users**

☒ Eppendorf tubes 1.5 mL uncolored **Eppendorf Centrifuge Catalog #022363204**

☒ MiniVortexer **VWR Scientific Catalog #58816-121**

☒ Surgical mask **Contributed by users**

☒ Micropipette (0.5 - 10 µL) **Contributed by users**

☒ Micropipette (100 - 1000 µL) **Contributed by users**

☒ pH meter **Contributed by users**

☒ Centrifuge 5424 R **Eppendorf Catalog #5404000022**

☒ Synergy™ HTX Multi-Mode Microplate Reader **Biotek**

STEP MATERIALS

☒ 55'-Dithiobis(2-nitrobenzoic acid) **Merck MilliporeSigma (Sigma-Aldrich) Catalog #D8130**

☒ Ethanol **Merck Millipore (EMD Millipore) Catalog #100983**

☒ Ethanol **Merck Millipore (EMD Millipore) Catalog #100983**

☒ Monobasic potassium phosphate **NUCLEAR Catalog #318312**

☒ Potassium phosphate dibasic **Neon Catalog #11361**

☒ Trichloroacetic acid (TCA) **Merck MilliporeSigma (Sigma-Aldrich) Catalog #T6399**

SAFETY WARNINGS

Use personal protective equipment (including lab coat, masks, and gloves) when manipulating chemical and biological samples. Read the Safety Data Sheets of the reagents.

BEFORE START INSTRUCTIONS

This protocol was standardized at LAPCOM (Psychopharmacology and Behavior Laboratory at UFRGS) to assess biochemical parameters in zebrafish brain tissue. Protocols you should read before proceeding with this method:

CITATION

Adrieli Sachett, Matheus Gallas-Lopes, Radharani Benvenuti, Greicy M M Conterato, Ana Herrmann, Angelo Piato. How to prepare zebrafish brain tissue samples for biochemical assays.

LINK

<https://protocols.io/view/how-to-prepare-zebrafish-brain-tissue-samples-for-bjkdks6>

CITATION

Adrieli Sachett, Matheus Gallas-Lopes, Greicy M M Conterato, Radharani Benvenuti, Ana Herrmann, Angelo Piato. Optimized protein quantification protocol for zebrafish brain tissue (Bradford method).

LINK

<https://protocols.io/view/optimized-protein-quantification-protocol-for-zebrbjnfkmbn>

Preparing the reagents

- 1 The first step is to prepare the reagents to be used in the quantification of nonprotein sulfhydryl groups (NPSH);

1.1 5,5'-Dithiobis(2-nitrobenzoic acid) (DTNB) [M] 10 mM :

1.1.1 Weigh carefully 0.0396 g of DTNB in a piece of aluminum foil;

5,5'-Dithiobis(2-nitrobenzoic acid) **Merck MilliporeSigma (Sigma-Aldrich) Catalog #D8130**

1.1.2 Transfer the DTNB to a beaker of appropriate size;

1.1.3 Add 9 mL of absolute ethanol to the beaker to dissolve the salt;

Ethanol **Merck Millipore (EMD Millipore) Catalog #100983**

1.1.4 Transfer your solution to a 10 mL volumetric flask;

1.1.5 Using absolute ethanol, complete the solution's volume to reach 10 mL ;

⊗ Ethanol **Merck Millipore (EMD Millipore) Catalog #100983**

1.1.6 Store the solution in an amber flask of appropriate size covered with aluminum foil at 8 °C ;

1.2 Potassium phosphate buffer [M] 1 Molarity (M) :

1.2.1 Weigh 13.609 g of monobasic potassium phosphate (KH_2PO_4) in a beaker of appropriate size;

⊗ Monobasic potassium phosphate **NUCLEAR Catalog #318312**

1.2.2 Dissolve the salt with 90 mL of ultrapure water;

1.2.3 Transfer the solution to a 100 mL volumetric flask;

1.2.4 Using ultrapure water, complete the solution's volume to reach 100 mL ;

1.2.5 Weigh 17.418 g of dibasic potassium phosphate (K_2HPO_4) in a beaker of appropriate size;

⊗ Potassium phosphate dibasic **Neon Catalog #11361**

1.2.6 Dissolve the salt with 90 mL of ultrapure water;

1.2.7 Transfer the solution to a 100 mL volumetric flask;

1.2.8 Mix both solutions slowly in a 500 mL beaker following the steps below;

- Transfer 50 mL of the monobasic potassium phosphate (KH_2PO_4) solution to the beaker;

- Use a pH sensor to evaluate your solution. Expected conditions: pH 7.0 ;

If the pH of your solution is lower than 7 adjust the pH adding drops of the dibasic potassium phosphate (K_2HPO_4) solution;

If the pH of your solution is above 7 adjust the pH adding drops of the monobasic potassium phosphate (KH_2PO_4) solution;

- After adjusting the pH of this initial solution, proceed to add, slowly, drops of both buffer solutions (monobasic potassium phosphate and dibasic potassium phosphate). Use Pasteur pipettes to add the solutions. Mix your solutions using a pH sensor, making sure the mix of both buffers is always at pH 7.0 ;

1.3 Trichloroacetic acid (TCA) 6%:

1.3.1 Weigh 6 g of TCA in a beaker of an appropriate size;

Trichloroacetic acid (TCA) **Merck MilliporeSigma (Sigma-Aldrich) Catalog #T6399**

1.3.2 Dissolve the TCA with 50 mL of ultrapure water;

1.3.3 Transfer your solution to a 100 mL volumetric flask;

1.3.4 Using ultrapure water, complete the solution's volume to reach 100 mL ;

1.3.5 Store this solution in an amber flask at 8 °C ;

Deproteinization of your samples

- 2 To proceed with the quantification of nonprotein sulfhydryl groups in your samples, you first have to deproteinize them following the steps below. Tissue sample collection and preparation are described elsewhere;

CITATION

Adrieli Sachett, Matheus Gallas-Lopes, Radharani Benvenuti, Greicy M M Conterato, Ana Herrmann, Angelo Piato. How to prepare zebrafish brain tissue samples for biochemical assays.

LINK

<https://protocols.io/view/how-to-prepare-zebrafish-brain-tissue-samples-for-bjkdks6>

- 2.1 Prepare 1.5 mL microtubes, to be used to store the samples, with the correct information. The number of microtubes depends on the number of samples;

- 2.2 Before preparing your samples for deproteinization, you must calculate the sample volume that corresponds to 50 µg of proteins. This calculation can be based on the Bradford method described elsewhere;

CITATION

Adrieli Sachett, Matheus Gallas-Lopes, Greicy M M Conterato, Radharani Benvenuti, Ana Herrmann, Angelo Piato. Optimized protein quantification protocol for zebrafish brain tissue (Bradford method).

LINK

<https://protocols.io/view/optimized-protein-quantification-protocol-for-zebr-bjnfkmbn>

2.2.1 To estimate the volume of the sample corresponding to 50 µg of proteins, divide the amount of protein needed (50 µg) by the total amount of proteins in the sample quantified by the Bradford method (example below);

Volume of the sample needed for the assay (µL) = 50 µg / total amount of proteins in the sample µg/µL

2.3 For each tissue sample, fill the plastic microtubes as described below. You should provide duplicates or triplicates of each sample to make your quantification more precise. The sample volume corresponding to 50 µg of proteins is the volume needed to fill one of the wells of the microplate, so if you are planning to evaluate your samples in duplicates fill the microtube with two times that volume, if evaluating in triplicates, three times that volume, and so on. Using a micropipette fill the tubes in this order: Sample and TCA solution (mixing the solution with the pipette tip to homogenize the content);

Sample (µL)	TCA 6% (µL)
Depends on the volume of the sample corresponding to 50 µg of proteins and the number of replicates of the same sample you are planning to evaluate	Depends on the volume of the sample. You should add the same volume of TCA solution as the volume of the sample you are planning to use (1:1)

2.4 Use a vortexer to mix the samples for 00:00:10 ;

2.5 Centrifuge the samples 10000 x g, 4°C, 00:05:00 ;

Microplate preparation and absorbance reading

3 Use a conventional 96-well microplate to run your samples. Reagents should be at room temperature. Pipetting of DTNB should be performed under dim or no light, making sure the microplate is carefully covered in aluminum foil to avoid photodegradation of the reagent;

3.1 Before start pipetting, each well of the microplate should be marked for sample identification;

3.2 Using an adequate micropipette, fill the wells of your microplate as described below. You should provide duplicates or triplicates of each sample as stated above. Using a micropipette fill the wells in this order: sample and TFK. The TFK volume depends on the volume of the sample. All wells should have a final volume of 245 μL , so the TFK is used so that every solution reaches this volume (e.g., 50 μL of the sample + 195 μL of the TFK solution);

Well of the plate	Sample (μL)	TFK 1 M (μL)
Control	0	245

Samples	Depends on the volume of the sample corresponding to 50 µg of proteins.	Depends on the volume of the sample. Volume needed for the final solution in the well to reach 245 µL.
---------	---	--

3.3 Read the absorbance of the samples at 412 nm in a microplate reader;

3.4 After reading the absorbance of the samples, add, in the dark, 15 µL of the DTNB solution [M] 10 mM to each well (control and samples) previously filled (mixing the solution with the pipette tip to homogenize the content of wells). The final volume of every well should reach 260 µL ;

3.5 Leave your microplate in a dark room to incubate at room temperature for 01:00:00 ;

3.6 Read the absorbance of the samples at 412 nm in a microplate reader;

Calculating data and determining results

4 Prepare to analyze the results obtained after reading the absorbance of the samples;

- 4.1 Calculate the mean absorbance of the control solution both before and after adding the DTNB;

$$\Delta_{\text{control}} = (\text{Mean absorbance after DTNB} - \text{Mean absorbance before DTNB})$$

- 4.2 Calculate the mean absorbance of the samples both before and after adding the DTNB;

$$\Delta_{\text{sample}} = (\text{Mean absorbance after DTNB} - \text{Mean absorbance before DTNB})$$

- 4.3 Subtract the Δ_{control} value from the Δ_{sample} value for each of the samples;

$$\text{Absorbance of the sample} = (\Delta_{\text{sample}} - \Delta_{\text{control}})$$

- 4.4 Determine the number of moles of the group sulfhydryl (SH) in the sample using the calculation below;

$$\text{Moles of SH} = \frac{\text{Abs}_{\text{sample}} \times 0.00021}{14.15 \times 0.67}$$

- 4.5 Results should be expressed as μmol of SH groups;

$$\mu\text{mol SH} = \text{Moles SH} \times 1000000$$

- 4.6 Calculate the amount of sulfhydryl groups per milligrams of protein;

$$\mu\text{mol SH/mg protein} = \frac{\mu\text{mol SH} \times 1000}{(\text{amount of proteins in the sample [50 } \mu\text{g in this case]})}$$

- 4.7 Final results are expressed as $\mu\text{mol SH/mg protein}$.