


May 14, 2024

Version 2

Quantitative analyses of the ultrastructural features of dopaminergic axon terminals V.2

DOI

<https://dx.doi.org/10.17504/protocols.io.bp2l694xdlqe/v2>



Natalie Doig¹, Peter Magill^{1,2}

¹Medical Research Council Brain Network Dynamics Unit, Nuffield Department of Clinical Neurosciences, University of Oxford, Mansfield Road, Oxford, OX1 3TH, United Kingdom;

²Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD



Cláudia C. Mendes

University of Oxford

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.bp2l694xdlqe/v2>

External link: <https://www.mrcbndu.ox.ac.uk/>

Collection Citation: Natalie Doig, Peter Magill 2024. Quantitative analyses of the ultrastructural features of dopaminergic axon terminals. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bp2l694xdlqe/v2> Version created by **Cláudia C. Mendes**



License: This is an open access collection distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this collection and it's working

Created: September 07, 2023

Last Modified: May 14, 2024

Collection Integer ID: 87520

Keywords: Dopamine, Axon terminal, Striatum Tyrosine hydroxylase (TH), Transmission Electron Microscopy (TEM), Electron microscopy (EM), Substantia nigra pars compacta (SNc), Ventral Tegmental Area (VTA), Vesicle Synapse, Ultrastructure, Immuno-EM, Immunogold, Quantification, ImageJ, user process animal brain tissue for electron microscopy, marker of dopaminergic axon, dopaminergic axon terminals in the brain, tissue sections ready for electron microscope, dopaminergic axon, dopamine neuron axon, dopaminergic axon terminal, electron microscopy, electron microscope, user process animal brain tissue, transmission electron microscope, image ultrathin sections of tissue, microscope, brain tissue, axon, release of dopamine, quantitative analyses of the ultrastructural feature, tissue section, dopamine, ultrastructural feature, inappropriate dopamine release, tissue

Abstract

The release of dopamine from axons is critical for normative brain function and behaviour. Impaired or otherwise inappropriate dopamine release often correlates with changes in the ultrastructure of dopamine neuron axons that can be assessed with electron microscopy. Here, we provide two protocols that can be used serially to, first, help the user process animal brain tissue for electron microscopy and, secondly, help the user undertake quantitative analyses of the ultrastructural features of dopaminergic axon terminals in the brain.

Protocol #1 describes how to prepare brain tissue, carry out pre-embedding immunohistochemistry for tyrosine hydroxylase as a marker of dopaminergic axons, and then make tissue sections ready for electron microscope.

Protocol #2 details how to examine and image ultrathin sections of tissue using a transmission electron microscope and then how to analyse the digital images.

Files

 SEARCH

Protocol



NAME

Quantitative analyses of the ultrastructural features of dopaminergic axon terminals. Protocol #1:
Tissue preparation for electron microscopy

VERSION 2

CREATED BY



Cláudia C. Mendes
University of Oxford

OPEN →

Protocol



NAME

Quantitative analyses of the ultrastructural features of dopaminergic axon terminals. Protocol #2:
Acquisition and analysis of electron microscopy images

VERSION 1

CREATED BY



Cláudia C. Mendes
University of Oxford

OPEN →