AUG 07, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.3bagiie

External link:
<https://www.diacomp.org/shared/document.aspx?id=222&docType=Protocol>

Protocol Citation: Sandra Rieger 2019. Quantitative PCR analysis to assess gene expression changes in hyperglycemic larval zebrafish. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.3bagiie>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: May 23, 2019

Last Modified: Aug 07, 2019

PROTOCOL integer ID:
 23618

Quantitative PCR analysis to assess gene expression changes in hyperglycemic larval zebrafish

Sandra Rieger¹

¹University of Miami

Diabetic Complications Consortium
 Tech. support email: rmcindoe@augusta.edu

Lili Liang

ABSTRACT

Summary:

To assess hyperglycemia in larval zebrafish, we analyzed the expression of genes implicated in glucose metabolism, such as insulin, insulin receptor, glucagon, and phosphoenolpyruvate carboxylase (pepck), using quantitative PCR (qPCR). We analyzed gene expression following treatment of Tg(*ins:NTRmCherry*) transgenic larvae either with 0.5 % DMSO (controls) or 10 mM metronidazole (see Protocol 2) for β -cell ablation at 3 and 8 days post fertilization (dpf).

Diabetic Complication:

Neuropathy

Keywords: PCR analysis, gene expression changes in hyperglycemic larval zebrafish, Neuropathy

MATERIALS

MATERIALS

✕ RNeasy Mini Kit **Qiagen Catalog #74104**

✕ oligo (dT) or random hexamer primers **Integrated DNA Technology**

✕ SuperScript®III First-Strand Synthesis System for RT-PCR **Life Technologies Catalog #18080-051**

✕ SYBR Green Brilliant III Ultra-Fast SYBR® Green QPCR Master Mix **Agilent Technologies Catalog #600882**

✕ ROX reference dye (included in Brilliant III master mix) **Agilent Technologies Catalog #600882**

✕ Mx3000P QPCR System **Stratagene Catalog #Mx3000P**

Primers:

Gene name	Gene symbol	5'-3' forward primer	5'-3' reverse primer
<i>insulin</i>	<i>ins</i>	c c c t t t a t c t g g t c t g t g g c	t t c e t t a t c a g c t c g g c a t g
phosphoenolpyruvate carboxylase	<i>pepck</i>	g a g a a c a g c a c c a t c e t c a g	t c a c e g t t t t a c t c t c c a c a c
<i>glucagon a</i>	<i>gcga</i>	a a g a c t c g t t c a g t g g c t c	t g g g t t g t c c g g a t t t a g c
<i>insulin receptor a</i>	<i>insra</i>	a t a a g a a t g a t c g g g a g t g t g g	t g c t t g c a t g a g g a t g g a c
<i>Elongation factor alpha (control gene)</i>	<i>efla</i>	t c t a c a a a t g c g g t g g a a t c g	g a g c a a t g t c a a t g g t g a t a c c

Note:

QIAGEN ([RRID:SCR_008539](#))

- 1 Preparation of mRNA from larval zebrafish. Pool 10 zebrafish larvae and isolate RNA according to the RNeasy Mini Kit (QIAGEN) manual. Elute final mRNA in 30 µl of RNase-free water.
- 2 Preparation of cDNA from mRNA. Prepare cDNA from isolated mRNA according to the SuperScript®III First-Strand Synthesis System (Life Technologies) using either oligo(dT) or random hexamer primers.
- 3 Preparation of qPCR mix (according to the Agilent manual for preparation of the Brilliant III Ultra-Fast SYBR® Green QPCR Master Mix):

Dilute the reference dye 1:500 using nuclease-free PCR-grade water. Prepare the experimental reactions by combining the components of the reagent mixture in the order listed in the table below. Prepare a single reagent mixture for replicate reactions (plus at least one reaction volume

excess) using multiples of each component.

Reagent Mixture

Nuclease-free PCR-grade water to bring final volume to 20 μ l (including cDNA)

10 μ l of 2 $\times$ SYBR Green QPCR Master Mix

x μ l of upstream primer at optimized concentration (200–500 nM)

x μ l of downstream primer at optimized concentration (200–500 nM)

0.3 μ l of diluted ROX reference dye

Gently mix the reagent mixture without creating bubbles, then distribute the mixture to the experimental reaction tubes. Add 0.5-50 ng (x μ l) of cDNA to each reaction to bring the final reaction volume to 20 μ l.

We amplified each transcript using the following PCR conditions:

1) 95°C 3 minutes

2) **40 cycles**

95°C 15 seconds

60°C 20 seconds,

3) **Melting curve**

95°C 1 minute

60°C 30 seconds

95°C 30 seconds

Potential Pitfalls:

- 4 Inconsistent qPCR results: Always use master mixes and the suggested reference dye (ROX reference dye, Life Technologies, USA). Include pre-mix of triplicate reactions with sample and primers. Alternatively, utilize the Taqman system (Life Technologies).