

Nov 17, 2020

Quantitative real-time PCR for the four Lactobacillus species

 [GigaByte](#)

✓ Peer-reviewed method

 In 4 collections

DOI

dx.doi.org/10.17504/protocols.io.bps2mnge

Chen Chen¹

¹BGI-SHENZHEN

GigaScience Press

BGI



Lilan Hao

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.bps2mnge

External link: <https://doi.org/10.46471/gigabyte.9>

Protocol Citation: Chen Chen 2020. Quantitative real-time PCR for the four Lactobacillus species. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bps2mnge>

Manuscript citation:

Chen C, Hao L, Wei W, Li F, Song L, Zhang X, Dai J, Jie Z, Li J, Song X, Wang Z, Zhang Z, Zeng L, Du H, Tang H, Zhang T, Yang H, Wang J, Brix S, Kristiansen K, Xu X, Wu R, Jia H, The female urinary microbiota in relation to the reproductive tract microbiota. GigaByte. 2020.

<https://doi.org/10.46471/gigabyte.9>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: November 17, 2020

Last Modified: November 24, 2020

Protocol Integer ID: 44602

Abstract

Quantitative Real-time Polymerase Chain Reaction(qPCR) for the four *Lactobacillus* species, including *L.crispatus*, *L.gasseri*, *L.iners*, *L.jensenii*.

- 1 The primers for the four *Lactobacillus* species used in this study were performed as Backer et al. (2007). Pick some vaginal DNA samples at random, and perform PCR reactions using these four sets of primers, respectively. Then use electrophoresis to identify the PCR products, and purify the final PCR product by gel extraction kit.

Specificity	Name	Primer sequence (5'-3')	16S rDNA position* (5'-3')	Cycling conditions
<i>L. crispatus</i>	LcrisF	AGCGAGCGGAACAAACAGATTTAC	65–89	10' 95°C, (15" 95°C, 1' 60°C) × 40
	LcrisR	AGCTGATCATGCGATCTGCTT	205–185	
<i>L. gasseri</i>	LactoF	TGGAAACAGRTGCTAATACCG	157–177	10' 95°C, (15" 95°C, 1' 60°C) × 40
	LgassR	CAGTTACTACCTCTATCTTTCTTCACTAC	470–442	
<i>L. iners</i>	InersFw	GTCTGCCTTGAAGATCGG	70–85	10' 95°C, (1' 95°C, 1' 55°C, 1' 65°C) × 35
	InersRev	ACAGTTGATAGGCATCATC	228–210	
<i>L. jensenii</i>	LABJENTR2Fw	CCTTAAGTCTGGGATACCATT	117–137	10' 95°C, (15" 95°C, 10" 54°C, 30" 72°C) × 40
	LABJENRT2Rev	ACGCCGCCTTTTAAACTTCTT	207–187	

- 2 Use the pEASYR-T1 Cloning Kit (https://www.transgenbiotech.com/cloning_vector.html) to construct the four corresponding plasmids and identify them by sanger sequencing, respectively. After DNA quantification using Qubit Fluorometer (Life Technologies), dilute the plasmids with ddH₂O by serial tenfold dilution into 6 concentration gradients as serial standards.
The copies of standards (copies/μl) = (DNA concentration/ (number of base pairs*324))*6*10¹⁴. The number of plasmid base pairs (bp) of *L. iners*, *L. jensenii*, *L. crispatus* and *L. gasseri* are 4082, 4013, 4082, 4251.
- 3 Use SYBR Premix Ex Taq GC (TAKARA) to conduct the qPCR. The reaction mixture contains 10 μl of 2×SYBR Premix Ex Taq GC, 0.2 μM forward primer, 0.2 μM reverse primer, 1.6 μl of DNA sample and 8.2 μl ddH₂O to make up the final reaction volume of 20 μl. Each reaction is run in triplicate.
- 4 Then transfer all the final reaction volumes into a 96-well plate. The DNA templates in each plate contain urine samples, corresponding serial standards and negative controls (ddH₂O).

- 5 Use the StepOnePlus Real-time PCR System (Life Technologies) to conduct the qPCR. The condition of qPCR amplification is followed as Table above. The standard curve range of amplification efficiency for the qPCR is from 90 to 110%, and linearity values is ≥ 0.99 . In case the result was not in the range of the standard curve, the samples are diluted tenfold and analyzed in triplicate again. The average log₁₀ copies/ml are expressed as per 1 ml urine sample.