VERSION 2
MAY 14, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.p26dqhe

External link:
<https://www.neb.com/protocols/2017/11/28/quick-protocol-for-monarch-total-rna-miniprep-kit-neb-t2010>

Protocol Citation: New England Biolabs 2018. Quick Protocol for Monarch® Total RNA Miniprep Kit (NEB #T2010). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.p26dqhe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: May 10, 2018

Last Modified: May 14, 2021

Quick Protocol for Monarch® Total RNA Miniprep Kit (NEB #T2010) V.2

New England Biolabs¹

¹New England Biolabs

New England Biolabs (NEB)

Tech. support phone: +1(800)632-7799 email: info@neb.com

Isabel Gautreau
New England Biolabs

ABSTRACT

Quick Protocol for Monarch® Total RNA Miniprep Kit ([NEB #T2010](#)).

Quickly and easily purify up to 100 µg of high-quality total RNA from multiple sample types – all with one kit!

- For use with blood, cells and tissues
- Also works with tough to lyse samples (bacteria, yeast, plant)
- Effectively purifies total RNA of all sizes, including small RNAs >20 nt
- Efficient genomic DNA removal (column and DNase I-based)
- Contains Proteinase K for processing of tissues and blood samples
- Includes RNA Protection Reagent for sample preservation
- Excellent value
- Kit components available separately

ATTACHMENTS

[T2010_Monarch_RNA_Kit_manualT2010.pdf](#)
[072817.pdf](#)

GUIDELINES

We recommend that first-time users of this kit review [the product manual](#) before starting. The manual provides additional relevant information to consider at various steps. This quick protocol is meant for experienced users.

RNA Purification Consists of Two Stages:

PART 1: Sample Disruption and Homogenization (Differs depending on starting material)

PART 2: RNA Binding and Elution

Have any questions?

Our tech support scientists would be happy to help. Email us at info@neb.com.

Keywords: total rna, rna extraction, rna purification, rna isolation

MATERIALS

MATERIALS

Monarch Total RNA Miniprep Kit **New England Biolabs Catalog #T2010S**

SAFETY WARNINGS

Please refer to the SDS for safety warnings.

BEFORE START INSTRUCTIONS

- Addition of RNA Lysis Buffer and all subsequent steps should be performed at room temperature to prevent formation of precipitate. If samples are accidentally placed on ice and precipitate forms, allow the samples to return to room temperature to resolubilize before loading onto the column.
- Protocols are also available for [RNA reaction cleanup](#), [RNA fractionation](#), and [extraction of RNA from other sample types](#) (including those in preservation reagents or TRIzol®).

For the 50 prep kit:

- Reconstitute DNase I by adding 275 µl nuclease-free water. Gently invert to mix. Aliquot for storage at -20°C to minimize freeze-thaw cycles.
- Reconstitute Proteinase K (Prot K) by adding 1040 µl of Proteinase K Resuspension Buffer. Vortex and store at -20°C.
- Add 100 ml ethanol (≥ 95%) to the 25 ml RNA Wash Buffer concentrate.

PART 1: Sample Disruption and Homogenization

- 1 Please select your starting material of the following:
 - **Cultured Mammalian Cells**
 - **Mammalian Whole Blood (Fresh or Frozen)**
 - **Tissue or Leukocytes**
 - **Tough-to-Lyse Samples (bacteria, yeast, plant, etc.) using Mechanical Lysis**

Step 1 includes a Step case.

Cultured Mammalian Cells

Mammalian Whole Blood (Fresh or Frozen)

Tissue or Leukocytes

Tough-to-Lyse Samples

step case

Cultured Mammalian Cells

- 2 Pellet cells by centrifugation (500 x *g*) for 1 min.

00:01:00 Centrifugation

- 3 Discard supernatant.

- 4 Resuspend pellet in RNA Lysis Buffer (according to table below) by pipetting gently to avoid foaming.

AMOUNT OF CELLS	VOLUME OF RNA LYSIS BUFFER
up to 3 x 10 ⁶	300 µl
3 x 10 ⁶ to 1 x 10 ⁷	≥ 600 µl

Note

Do not place samples on ice. For frozen pellets, thaw briefly before resuspension.

PART 2: RNA BINDING AND ELUTION

- 5 Transfer up to 800 µl of the sample from PART 1 to a gDNA removal column

(light blue) fitted with a collection tube.

For sample identification, **label collection tubes**, as gDNA removal columns will be discarded after spinning.

Note

For sample volumes > 800 µl (column reservoir capacity), columns may be reloaded.

Note

*All centrifugation steps should be performed at 16,000 x *g*.*

- 6 Spin for 30 seconds to remove most of the gDNA. **SAVE THE FLOW-THROUGH** (RNA partitions here). Discard the gDNA removal column.

00:00:30 Spinning

- 7 Add an equal volume of ethanol (≥ 95%) to the flow-through and mix thoroughly by pipetting. **Do**

not vortex. To exclude RNA ≥ 200 nt, add only 1/2 volume ethanol to flow-through.

8 Transfer mixture to an RNA purification column

(dark blue) fitted with a collection tube. Spin for 30 seconds. Discard flow-through. If further gDNA removal is essential for downstream applications, proceed to on-column DNase I treatment, steps 9-11 (recommended). If not, proceed to **Step 12**.

00:00:30 Spinning

9 **[Optional (but recommended)]** On-column DNase I treatment for enzymatic removal of residual gDNA:

Add 500 μ l RNA Wash Buffer and spin for 30 seconds. Discard flow-through.

500 μ L RNA Wash Buffer

00:00:30 Spinning

10 **[Optional (but recommended)]** In an RNase-free microfuge tube (not included), combine 5 μ l DNase I with 75 μ l DNase I Reaction Buffer and pipet directly to the top of the column matrix.

5 μ L DNase I

75 μ L DNase I Reaction Buffer

11 **[Optional (but recommended)]** Incubate for 15 minutes at room temperature.

00:15:00 Incubation

12 Add 500 μ l RNA Priming Buffer and spin for 30 seconds. Discard flow-through.

500 μ L RNA Priming Buffer

00:00:30 Spinning

13 Add 500 μ l RNA Wash Buffer and spin for 30 seconds. Discard flow-through.

500 μ L RNA Wash Buffer

00:00:30 Spinning

14 Add another 500 μ l RNA Wash Buffer and spin for 2 MINUTES.

500 μ L RNA Wash Buffer

00:02:00 Spinning

- 15** Transfer column to an RNase-free microfuge tube (not included). Use care to ensure the tip of the column does not contact the flow-through. If in doubt, re-spin for 1 minute to ensure no ethanol is carried over.

00:01:00 Re-spinning

- 16** Add 30-100 µl Nuclease-free Water directly to the center of column matrix and spin for 30 seconds.

00:00:30 Spinning

Note

For best results, elute with at least 50 µl, which is the minimum volume needed to wet the membrane. Lower volumes can be used but will result in lower recovery (elution in 30 µl results in > 80% recovery and 100 µl provides maximum recovery). For spectrophotometric analysis of eluted RNA, it may be necessary to re-spin eluted samples and pipet aliquot from top of the liquid to ensure that the $A_{260/230}$ is unaffected by possible elution of silica particles.

- 17** Place RNA on ice if being used for downstream steps, at **-20°C** for **short-term storage** (less than 1 week), or at **-80°C** for **long-term storage**. Addition of EDTA to 0.1-1.0 mM may reduce the activity of any contaminating RNases.