APR 24, 2019

🌐 Quick staining procedure of nuclei in Euplotes using DAPI

🔗 Forked from [Quick staining procedure of nuclei in Euplotes using DAPI](#)

📁 In 1 collection

Rachele Cesaroni¹

¹Universität Bern

Protist Research to Optimize Tools in Genetics (PROT-G)

OPEN ACCESS

Angela Piersanti
University of Camerino

DOI:
dx.doi.org/10.17504/protocols.io.2ajgacn

Protocol Citation: Rachele Cesaroni 2019. Quick staining procedure of nuclei in Euplotes using DAPI. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.2ajgacn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Apr 24, 2019

Last Modified: Apr 24, 2019

PROTOCOL integer ID:
22571

- 1 Mix concentrated Euplotes cells together with Ethanol 70% in a ratio of 1:1.

Note

Better to have a completely starved Euplotes culture to avoid autofluorescence from bacteria/algae.

- 2 Add DAPI (0.01 mg/ml) to the mix in a ratio of 1:10, and stain for 15 minutes at room temperature.

- 3 Observe cells by fluorescence microscopy.