



Jan 05, 2024

Version 1

Rat organotypic cultures for AAV-mediated vital labeling of the extracellular matrix V.1

DOI

dx.doi.org/10.17504/protocols.io.e6nvwdnzzlmk/v1

Federico N. Soria¹, Mario Fernandez-Ballester¹

¹Achucarro Basque Center for Neuroscience



Federico N. Soria

Achucarro Basque Center for Neuroscience

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.e6nvwdnzzlmk/v1>

Protocol Citation: Federico N. Soria, Mario Fernandez-Ballester 2024. Rat organotypic cultures for AAV-mediated vital labeling of the extracellular matrix. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.e6nvwdnzzlmk/v1>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working



Created: January 05, 2024

Last Modified: May 31, 2024

Protocol Integer ID: 92986

Keywords: ASAPCRN, rat organotypic cultures for aav, vital labeling of the extracellular matrix, imaging of the extracellular matrix, exportable hyaluronan, extracellular matrix, organotypic culture, neurocan, mediated vital labeling, aav, vital labeling

Funders Acknowledgements:

Aligning Science Across Parkinson









Grant ID: ASAP-020505

Abstract


In this protocol we describe the preparation and maintenance of rat organotypic cultures (**cortico-striatal**) for AAV-infection and imaging of the extracellular matrix. The AAV, in this case, induces the expression of a exportable hyaluronan-binding protein (Neurocan) fused with GFP, that allows vital labeling (and hence live imaging) of the extracellular matrix. The organotypic cultures can be maintained for several weeks, allowing for repeated imaging and longitudinal studies.

Materials


Organotypic culture medium

- 50%  MEM, HEPES, no glutamine **Gibco - Thermo Fisher Scientific Catalog #32360026**
- 25%  EBSS, calcium, magnesium, phenol red **Thermo Fisher Catalog #24010043**
- 25%  Horse Serum, heat inactivated, New Zealand origin **Thermo Fisher Catalog #26050088**
- 1x  B-27 Supplement **Gibco - Thermo Fisher Scientific Catalog #17504044**
- 1x  GlutaMAX™ Supplement **Thermo Fisher Scientific Catalog #35050061**
- 36 mM  Glucose Solution **Thermo Fisher Catalog #A2494001**
- 50 U/mL  Penicillin-Streptomycin (5,000 U/mL) **Thermo Fisher Catalog #15070063**
- 1.25 ug/mL  Fungizone (Amphotericin B) **Thermo Fisher Scientific Catalog #15290018**

Membrane Inserts

-  Millipore Cell Culture Inserts (0.4 um pore size, 30 mm diameter) **Merck MilliporeSigma (Sigma-Aldrich) Catalog #PICM0RG50**

Tissue Chopper

Equipment		
McIlwain Tissue Chopper	NAME	
Cavey Laboratory Engineering	BRAND	
51350	SKU	
		




Protocol materials

- ✕ HBSS, no calcium, no magnesium, no phenol red **Thermo Fisher Scientific Catalog #14175095**
- ✕ Horse Serum, heat inactivated, New Zealand origin **Thermo Fisher Catalog #26050088**
- ✕ B-27 Supplement **Gibco - Thermo Fisher Scientific Catalog #17504044**
- ✕ GlutaMAX™ Supplement **Thermo Fisher Scientific Catalog #35050061**
- ✕ Millipore Cell Culture Inserts (0.4 um pore size, 30 mm diameter) **Merck MilliporeSigma (Sigma-Aldrich) Catalog #PICM0RG50**
- ✕ MEM, HEPES, no glutamine **Gibco - Thermo Fisher Scientific Catalog #32360026**
- ✕ EBSS, calcium, magnesium, phenol red **Thermo Fisher Catalog #24010043**
- ✕ Glucose Solution **Thermo Fisher Catalog #A2494001**
- ✕ Penicillin-Streptomycin (5,000 U/mL) **Thermo Fisher Catalog #15070063**
- ✕ Fungizone (Amphotericin B) **Thermo Fisher Scientific Catalog #15290018**
- ✕ Cytosine β -D-arabinofuranoside (AraC) **Merck MilliporeSigma (Sigma-Aldrich) Catalog #C1768**

Troubleshooting



Culture preparation

- 1 Prepare a 6-well culture plate with Millipore 0.4 μm culture inserts placed on 1 mL of organotypic medium per well, and incubate it at  37 °C and 5% CO₂ to warm up.


Note


It is important to minimize air bubbles between the medium and the membrane insert, as this will prevent access of nutrients to the slice.

- 2 Prepare a p100 and p60 petri dishes with 25 mL and 5 mL, respectively, of



HBSS, no calcium, no magnesium, no phenol red **Thermo Fisher Scientific Catalog #14175095**

and keep them  On ice .

- 3 Prepare the **McIlwain Tissue Chopper** (see Materials) with a fresh blade and clean holder plates.
- 4 Sacrifice P5-P7 rat pups by rapid decapitation and extract the brain without damaging the cortex.
- 5 Put the brain in the on a filter paper and cut through the midline to separate the cerebral hemispheres. Remove the midbrain, separating it from the cortex with a scalpel.
- 6 Place both hemispheres in the same holder plate, with the medial face downward. Hold the plate with both hemispheres perpendicular to the blade (coronal sections) and section the brain at 350 μm width.
- 7 Pour the sliced brain carefully into the p100 petri dish with HBSS  On ice . You may use a Pasteur pipette to moist the brain and displace it more easily.
- 8 Separate slices under the magnification microscope with a couple of spatulas (preferentially plastic-ones, which can be custom crafted cutting the back of a Pasteur pipette). Use a blade and clean cuts to remove unwanted regions (such as hippocampus



or SVZ). Transfer the slices into the p60 petri dish (you can use an inverted glass pipette with a suction rubber bulb).


Note

It is important to remove the subventricular zone (SVZ) and the meninges with the scalpel to avoid cell proliferation.

- 9 **Under the laminar flow hood**, transfer the selected slices (undamaged slices from the preferred rostro-caudal levels) to preheated culture inserts with medium (Step 1). Each insert can only hold two cortico-striatal slices. Remove excess dissection buffer from the top of the insert with a sterile tip.

Note

Leaving too much HBSS solution on top of the insert will eventually dilute the feeding medium.

- 10 Incubate the culture at  37 °C and 5% CO₂.

Expected result

Typically, 4-6 cortico-striatal slices can be obtained from each hemisphere

Maintenance and AAV-infection

- 11 At DIV1 (24h after culture preparation), replace the feeding medium with 1 mL of the same medium but adding antimetabolic



Cytosine β -D-arabinofuranoside (AraC) Merck MilliporeSigma (Sigma-Aldrich) Catalog #C1768

4,4 μ m per well.



Expected result

AraC will prevent the overproliferation of astrocytes, minimizing the formation of a glial scar on top of the slice.

- 12 At DIV3, replace the AraC-containing medium with fresh medium. Then add 1 μ L of the virus of choice (in our case, AAV-Ncan-GFP 10^{12} vg/mL) over the slice.

Safety information

AAVs have to be manipulated in BSL1 facilities.

Note

To improve spatial precision (e.g. when infecting a selected region from the slice), AAV infection can be done with a micromanipulator and a glass pipette.

- 13 To maintain the culture, change medium three times a week (1 mL per well).

Expected result

Slices will flatten overtime, and become more transparent. Cortical regions will have more cell density than striatal regions.

AAV expression become visible aprox. at DIV10, with maximum expression from DIV14 onwards (it depends on AAV serotype and promoter).

Imaging

- 14 Imaging can be performed in two ways:
- **IN-PLATE:** The insert **is not removed from the plate**, and the imaging is performed through the plastic. This is usually done in *widefield* setups, with low magnification. It is not optimal in terms of resolution, but it is very useful for repeated imaging (longitudinal studies) without compromising sterility.
 - **OFF-PLATE:** The insert is **removed from the plate** and placed in a smaller petri-dish or similar holding chamber. Imaging is usually done from above, in upright microscopes (objective on-top), with water-based objectives. Since CO₂ is usually not available, a buffer medium with HEPES is recommended. Phenol-red should be



avoided as well, since it interferes with fluorescence. This type of configuration is typically used for *confocal* or *2-photon* imaging of the slices.