MAY 14, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.p4ndqve

Protocol Citation: Katy M Monteith 2018. Recipe for apple-agar embryo collection plates. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.p4ndqve>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: May 14, 2018

Last Modified: May 14, 2018

PROTOCOL integer ID:
 12142

Keywords: apple-agar, embryo collection

Recipe for apple-agar embryo collection plates

Katy M Monteith¹

¹University of Edinburgh

Vale Lab

Katy M Monteith
 University of Edinburgh

ABSTRACT

This protocol describes the recipe for apple-agar plates which can be used for embryo collection. Alternatively, grape juice can be used instead of apple juice.

GUIDELINES

This protocol describes how to make sufficient apple-agar for approximately 10x 90mm Petri dish plates.

Once the apple-agar food is decanted, the plates should be left, covered by lids, at room temperature until the food has set and cooled. The plates should then be sealed in an air tight bag and stored at 4°C for a maximum of ~ 6 weeks before use. The plates should be brought to room temperature before being added to a population cage.

MATERIALS

MATERIALS

- ☒ Sucrose **Sigma Aldrich Catalog #S7903**
- ☒ Agar ash 2.0-4.5% **Sigma Aldrich Catalog #A7002**
- ☒ Methyl-4- hydroxybenzoate **Sigma Aldrich Catalog #H5501**

SAFETY WARNINGS

1. Care should be taken when dispensing hot apple-agar liquid. Decant liquid into smaller container before pouring into plates to reduce the risk of spillage.
2. Read SDS for Methyl 4-hydroxybenzoate prior to use.

- 1 Add 9 g of agar and 10 g sucrose to 300 mL distilled H₂O in a pan.

- 2 Mix well, place pan on heat and bring mixture to boil while stirring continuously.
- 3 Once boiling, turn off heat and add 100 mL apple juice to mixture.
- 4 Allow mixture to cool to ~70 °C and then stir in 2.1 mL of tegosept anti-fungal solution (dissolve Methyl-4- hydroxybenzoate in 95% Ethanol at 150g/L).
- 5 Dispense apple-agar into plates before it sets.