

Oct 02, 2025

REDI-NET FV-3 FLYING VECTOR STORAGE

DOI

<https://dx.doi.org/10.17504/protocols.io.81wgbkm11gpk/v1>



REDI-NET Consortium¹

¹REDI-NET Consortium

Remote Emerging Disea...



REDI-NET Consortium

University of Notre Dame, Naval Medical Research Center, Wal...

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.81wgbkm11gpk/v1>

Protocol Citation: REDI-NET Consortium 2025. REDI-NET FV-3 FLYING VECTOR STORAGE. protocols.io <https://dx.doi.org/10.17504/protocols.io.81wgbkm11gpk/v1>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: June 20, 2025

Last Modified: October 02, 2025

Protocol Integer ID: 220633

Keywords: STORAGE PROCEDURE FOR UNTREATED SAMPLE, STORAGE PROCEDURE FOR TOTAL NUCLEIC ACID, flying vector, storage, flying vector storage, vector storage, vector, redi

Funders Acknowledgements:

USAMRAA

Grant ID: W81XWH-21-C-0001

USAMRAA

Grant ID: W81XWH-22-C-0093

USAMRAA

Grant ID: HT9425-23-C-0059

USAMRAA

Grant ID: HT9425-24-C-0072

Disclaimer

This work is supported by the US Army Medical Research and Development Command under Contract No.W81XWH-21-C-0001, W81XWH-22-C-0093, HT9425-23-C-0059 and HT9425-24-C-0072. The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army or Navy position, policy or decision unless so designated by other documentation.

Abstract

This protocol describes flying vector storage.



Guidelines

OBJECTIVE

To outline steps for properly storing blood samples and nucleic acid samples purified from these samples.

SUMMARY/SCOPE

The overarching aim of the REDI-NET is to develop a collaborative laboratory network between domestic and international partnering institutions to address disease surveillance needs in order to effectively detect, predict and contain potentially emergent zoonosis. This SOP provides guidance on storage of blood samples and their purified nucleic acid to preserve their integrity for downstream nucleic acid extraction and/or sequencing library preparation

MAINTENANCE OF EQUIPMENT

Decontaminate a PCR workstation by keeping the UV light on for  00:15:00 .

Materials

EQUIPMENT AND MATERIALS

Note

NOTE: If product number is listed, please ensure use of this or equivalent product.

A	B	C
Equipment / Material	Description	Mfg / Product #
-80°C freezer	For sample storage	Locally sourced
Forceps	Clean, stainless	Locally sourced
Ice	To maintain cold chain during sample handling	Locally sourced
96-Well Microfuge tube racks with cover	To hold microplates	Locally sourced
KingFisher™ 96 KF microplate	To store the sample	ThermoFisher, 97002540
PCR Workstation	PCR cabinet with UV light	Locally sourced
Clear Adhesive Film	To seal the KingFisher™ 96 KF microplate	ThermoFisher, 4306311
Adjustable micropipettes	To handle the samples	Locally sourced
Multi-channel micropipettes	8- or 12- channel; to handle the sample	Locally sourced
Nuclease-free filter tips low-retention	To ensure appropriate sample handling	Locally sourced
Nuclease free microfuge tubes	1.5 mL	Locally sourced
Saran wrap	Plastic wrap; to seal rack holding sample	Locally sourced
Permanent markers	To label tubes and microplates	Locally sourced
Data sheet	REDI-NET DCS T-3 Tick Storage	REDI-NET Data Portal



Safety warnings

RISKS AND PERSONAL PROTECTION:

Gloves should be worn all the time when handling samples.

Ethics statement

The protocols.io team notes that research involving animals and humans, including tissues or samples collected directly from specimens or participants, must be conducted according to internationally-accepted standards and should always have prior approval from an Institutional Ethics Committee or Board.



STORAGE PROCEDURE FOR UNTREATED SAMPLE

1

Note

NOTES:

- Flying vector samples need to be kept on a cold chain all the time to prevent RNA degradation. The following procedure will apply only where -80°C storage is feasible.
- If -80°C storage is not possible, temporarily store the flying vector samples in a -20 °C freezer and follow flying vector sample processing SOP (REDI-NET SOP FF-2 Flying Vector Processing) as soon as possible for total nucleic acid extraction. Subsequently, use a portion of the total nucleic acid and reverse-transcribe RNA into cDNA for -20 °C storage. To do this, follow the initial steps of the flying vector sample testing SOP (REDI-NET SOP FF-4 Flying Vector Testing) till finishing the step of "Removal of RNA".

Precool 96-well microfuge tube racks  On ice .

- 2 Using preprinted adhesive labels or permanent markers, label 1.5 ml microfuge tubes with unique sample ID.
- 3 Transfer flying vector sample into the corresponding pre-labeled 1.5 mL microfuge tubes, close the cap and put it onto the microfuge tube rack ( On ice) sequentially.
- 4 Once the rack is full or all flying vector samples have been completed, label the rack with a unique rack ID.
- 5 Close the rack lid tightly, secure with clear Saran wrap and immediately transfer to  -80 °C freezer.
- 6 Update the freezer inventory so samples can be tracked properly.

STORAGE PROCEDURE FOR TOTAL NUCLEIC ACID

7





Note

NOTES:

- The following procedure is to properly store total nucleic acid extracted from blood samples (including negative controls) using KingFisher nucleic acid purification system. The eluted total nucleic acid will be in either 96-well microplate (Flex model) or elution strip (Duo Prime model).
- Total nucleic acid samples need to be kept  On ice all the time to minimize RNA degradation.

In the clean PCR workstation, carefully transfer the eluted total nucleic acid to a 96-well PCR microplate, make sure to keep samples in the exact same locations corresponding to the rack where the original blood samples were stored.

Note

IMPORTANT: Mark the “A1” position of the 96-well microplate to prevent any mistakes on plate orientation.

- 8 Cover the 96-well PCR microplate with adhesive film to prevent spill over or contamination.
- 9 Label the film with a unique plate ID.
- 10 Immediately transfer the 96-well PCR microplate to  -80 °C freezer.
- 11 Update the freezer inventory so samples can be tracked properly.

Protocol references

REFERENCES

[REDI-NET Overview Summary](#)