MAR 06, 2020

Reducing Power

PLOS One

Jing Xu¹

¹Hainan University

Jing Xu

Hainan University

ABSTRACT

The reducing power of samples were assayed using the Guo et al. 1 ml of the extract was added to 2.5 ml of phosphate buffer (0.2 M, pH 6.6) and 2.5 ml of 1% potassium ferricyanide. After 20 min, 2.5 ml of 10% trichloroacetic acid (TCA) was added, followed by centrifugation at 3000 rpm for 10 min. The upper layer (2.5 ml) was mixed with 2.5 ml of distilled water and 0.5 ml of 0.1% ferric chloride, after 10 min, the absorbance was measured at 700 nm. EC₅₀ was calculated from the graph of an inhibition percentage against extract concentration.

OPEN ACCESS

DOI:

[dx.doi.org/10.17504/protocols.io.bda2i2ge](https://doi.org/10.17504/protocols.io.bda2i2ge)

External link:

<https://doi.org/10.1371/journal.pone.0234435>

Protocol Citation: Jing Xu 2020. Reducing Power.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bda2i2ge>

MANUSCRIPT CITATION:

Zhou J, Yang Q, Zhu X, Lin T, Hao D, Xu J (2020) Antioxidant activities of *Clerodendrum cyrtophyllum* Turcz leaf extracts and their major components. PLoS ONE 15(6): e0234435. doi: [10.1371/journal.pone.0234435](https://doi.org/10.1371/journal.pone.0234435)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Mar 06, 2020

Last Modified: Mar 06, 2020

PROTOCOL integer ID:
33850