SEP 22, 2019

Removal of kanamycinR gene from Keio collection strain

Ben Kuipers¹

¹Wageningen University

iGEM Wageningen 2019

Ben Kuipers

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.7jqhkmw

Protocol Citation: Ben Kuipers 2019. Removal of kanamycinR gene from Keio collection strain. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.7jqhkmw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol in our group and it is working. (This protocol can also be completed in 3 days, starting after the overnight culture).

Created: Sep 22, 2019

Last Modified: Sep 22, 2019

PROTOCOL integer ID:
 27984

ABSTRACT

This protocol is used to remove the kanamycinR gene from the JW3367 strain from the Keio Collection.

References Datsenko, K.A., and Wanner, B.L. (2000) One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. U.S.A.* 97: 6640-6645.

Barrick Lab. (16 May, 2014). FLP Recombination in *E. coli*. Retrieved 3 September, 2016, from <http://barricklab.org/twiki/bin/view/Lab/ProcedureFLPFRTRecombination>
 T-Hong_Kong_HKUST--Protocol_RemoveKAN.pdf

- 1 Chemical competent cells of *E. coli* JW0336 were prepared with the mix and go protocol.

- 2** Day 1: Transformation of the recombinase plasmid pCP20
- The competent cells were transformed with plasmid pCP20. (This plasmid has a temperature-sensitive origin of replication, resistant to ampicillin and chloramphenicol. With the presence of the FLP recombinase, the ampicillin and chloramphenicol resistance genes would be removed under the temperature higher than 43°C).
 - 1 ml LB was added for recovery which lasted for 1 hour.
 - The eppendorf was centrifuged for 2.5 minutes with 11000xg. Supernatant was discarded and the pellet was resuspended and plated on the LB + AMP plate.

The plate was then incubated for 1.5 days under 30°C.

- 3** Day 2: Induction of the recombination
- Three colonies were picked and inoculated into 5 mL liquid LB cultures.

The tubes were shaken overnight at 45°C so as to induce FLP recombinase expression and select for the loss of pCP20.

- 4** Day 3: Plating to get single candidate recombinants
- 100-fold dilution of the overnight culture was made by using fresh LB.
 - 50 ul of the diluted overnight culture was plated on LB plate.

The plate was then incubated for 1.5 days under 30°C to prevent partial loss of plasmid from the colonies which developed from the cells containing pCP20.

- 5** Day 4: Screening for genomic recombination and plasmid loss
- 6 colonies were picked from each plate and resuspended in MilliQ
 - Little amount of the MilliQ with cells was streaked on the LB plate(without antibiotics). The remaining solution was divided over LB + AMP, LB + CHL and LB + KAN plates dropwise for screening.

The plate was then incubated overnight under 37°C.

- 6** Day 5: Obtaining successful recombinants.
- The candidate which was sensitive to all antibiotics would be selected and archived.