AUG 13, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.skdecs6

Protocol Citation: Leonardo Fortes 2018. Resistance training volume protocol vs. HRV. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.skdecs6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 13, 2018

Last Modified: Aug 13, 2018

PROTOCOL integer ID:
 14693

Keywords: heart rate;
 resistance training;

Resistance training volume protocol vs. HRV

Leonardo Fortes¹

¹Universidade Federal da Paraíba

Leonardo Fortes

ABSTRACT

This is a controlled, randomized, and cross-over investigation with 40-week duration. The participants were male young adults with previous experience with RT for testing the effect of long-term RT with different volumes on HRV. This study was conducted in healthy participants in order to investigate the effects of resistance training on HRV without pathological influences. They underwent to a 40-week intervention divided in three RT segments (8 weeks each) and two washouts (8 weeks each) according Figure 1. The intervention order was randomized and counterbalanced. Thus, participants performed 1, 3 or 5 sets of the same exercises with equalized intensity (loading zones) and rest (Table 1) for 8 weeks following 8 weeks of washout between each experimental condition (1 vs. 3 vs. 5 sets). 10 repetition maximum (10-RM) test was performed for defining training intensity that was adjusted weekly (increase/decrease of 2-5% and 5-10% for upper and lower limbs, respectively) as recommended by Ribeiro et al. (2016). In case the participant performed more than 10 repetitions in two sessions, load (kg) was increased in the next week, whereas whether the participant was not able to perform 10-RM the load was decreased in the next week.

HRV was assessed by cardiac monitoring 72h before (pre) and after (post) each experimental RT condition (1 vs. 3 vs. 5 sets). Moreover, the participants were instructed to not performing any physical exercise 48h previously HRV baseline evaluation.

ATTACHMENTS

[Figura 1 - English.jpg](#)

GUIDELINES

American College of Sports Medicine

SAFETY WARNINGS

- ! For the training program, we follow previous recommendation for RT in healthy adults (ACSM, 2009). The program was composed by four exercises (bench press, leg press 45°, seated row, leg curl) performed three times per week (Monday, Wednesday, and Friday) during 8 weeks for each experimental approach (1 vs. 3 vs. 5 sets). All sessions were performed during the afternoon period (4 p.m.) to avoid circadian rhythm effect. All sessions were supervised by experienced researchers in RT area.

Table 1 shows RT procedure in each condition (1 vs. 3 vs. 5 sets). Warm-up (1 x 20-25 reps with 50% of predicted 10RM and 1 x 10-15 reps with 80% of predicted 10RM, between sets was given 3 min interval) was performed for the first two exercises (bench press and leg press 45°) before each resistance training session. Moreover, it is noteworthy to point out that during the washout period the participants did not perform RT.