

Jan 16, 2020

Version 1

Respiratory distress syndrome of the newborn and transient tachypnea of the newborn diagnosis V.1

DOI

dx.doi.org/10.17504/protocols.io.bavtie6n



Marconi Augusto Aguiar dos Reis¹, Roberta Maia C Romanelli¹, Zilma Reis¹

¹Universidade Federal de Minas Gerais



Zilma Reis

UFMG

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.bavtie6n>

Protocol Citation: Marconi Augusto Aguiar dos Reis, Roberta Maia C Romanelli, Zilma Reis 2020. Respiratory distress syndrome of the newborn and transient tachypnea of the newborn diagnosis . **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.bavtie6n>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Our multicenter team of researchers uses this protocol to recognize respiratory distress syndrome of newborn (SDR) and transient tachypnea of the newborn (TTN), secondary outcomes in the International Clinical Trials Registry Platform under the numbers RBR-3f5bm5 and RBR33rnjf.

Created: December 30, 2019

Last Modified: January 16, 2020

Protocol Integer ID: 31379

Keywords: Respiratory distress syndrome of newborn, transient tachypnea of newborn, respiratory disorders of newborn, prematurity, transient tachypnea of the newborn diagnosis, newborn diagnosis, transient tachypnea, respiratory distress syndrome, enrolled newborn, related respiratory disorder, newborn, respiratory disorders during the first 72h, test multicentre clinical trial in brazilian hospital, protocol for the preemy, low birth weight, international multicenter trial protocol for classification, diagnosis, new medical device, prematurity detection, accordance with the good clinical practice protocol, international clinical trials registry platform, clinical trial, good clinical practice protocol, standard procedures for immaturity

Abstract

The authors detailed describe the standard procedures for immaturity-related respiratory disorders during the first 72h of life. These disorders are respiratory distress syndrome of newborn (SDR) and transient tachypnea of the newborn (TTN), secondary outcomes in the International Clinical Trials Registry Platform under the numbers **RBR-3f5bm5** and **RBR33rnjf**.

It shall be used by the multicenter team of researchers, duly trained in accordance with the Good Clinical Practice Protocol, during the prospective evaluation of enrolled newborns. Also, this protocol is complementary documentation for the scientific publications related to the clinical trials:

"Prematurity detection evaluating interaction between the skin of the newborn and light: protocol for the preemie-test multicentre clinical trial in Brazilian hospitals to validate a new medical device".

"Premature or small for gestational age? International multicenter trial protocol for classification of the low birth weight newborn through the optical properties of the skin".



Guidelines

The scientific references for the diagnosis of respiratory distress syndrome of newborn (RDS) and transient tachypnea of the newborn (TTN) were:

- 1 - Liszewski MC, Stanescu AL, Phillips GS, et al. Respiratory Distress in Neonates: Underlying Causes and Current Imaging Assessment. *Radiol Clin North Am* 2017;55:629–44. 7.
- 2 - Reuter S, Moser C, Baack M. Respiratory distress in the newborn. *Pediatr Rev* 2014;35:417–29. 20.
- 3 - Silverman, W. A., & Andersen, D. H. (1956). A controlled clinical trial of effects of water mist on obstructive respiratory signs, death rate and necropsy findings among premature infants. *Pediatrics*, 17(1), 1-10.

Materials

Standardized Clinical Trial Data Collection Form and Tablet for recording data of the study

Troubleshooting

Safety warnings

- ! The newborns received treatment for RDS, TTN and other complications, according to the best clinical practice and with the local protocols, without any influence of this study.

Before start

Identify the sources of the hospital records during the first 72 hours of life: medical records, laboratory and radiological assessments of the newborn.

Respiratory distress syndrome of the newborn

1 The following clinical characteristics characterize this condition

- 1.1
 - a) Tachypnea that persists after 2 hours of life without respiratory distress or mild respiratory distress (maximum score of 3 - according to a clinical description by Silverman Anderson Bulletin (1956).
 - b) Self-limited tachypnea, worsening up to 24 hours and later improvement, with resolution within 72 hours of life or discharge from hospital.
 - c) Newborns with diagnosed extrapulmonary conditions who have tachypnea (cardiac, neurological, metabolic) will be excluded from the study.
- 1.2 Other complementary findings that may or may not be present during data collection were:
 - a) Chest X-ray showing diffuse interstitial infiltrates with distribution pattern from the hilum, lobar fissure thickening, pleural effusion (usually small) or opacity for perihilar reinforcement (in severe cases). There may be mild to moderate cardiomegaly.
 - b) Respiratory distress was assessed according to the Silver Andersen respiratory severity score (RSS) for both respiratory disorders

Transient tachypnea of the newborn

2 The following clinical characteristics characterize this condition

- 2.1
 - A. Tachypnea that persists after 2 hours of life with signs of progressive worsening respiratory distress
 - B. Tachypnea onset within the first 6 hours of life, worsening within 24 hours and requiring ventilatory support.
 - C. Absence of extrapulmonary conditions that cause tachypnea from other causes such as cardiac, neurological, metabolic, and infectious. However, when there is no perinatal asphyxia (due to these diagnoses).



- 2.2 Criteria A + B + C were considered mandatory for the diagnosis. Other complementary findings that may or may not be present:
- a) Use of surfactant in the first 24 hours of life.
 - b) Chest X-ray performed within 24 hours of life (or before surfactant) showing granular opacity or diffuse pulmonary reticulogranular infiltrate (ground-glass appearance) and reduced lung volume with air bronchogram (indicative of surfactant's deficiency).
 - c) Saturation below target saturation (92-95%) with progressive worsening and need for higher levels of FiO₂ (with FiO₂ above 35 to 45 to maintain PO₂ > 50mmHg - blood gas revealing respiratory acidosis)

Diagnosis confirmation by the experts

- 3 Two experts in neonatology reviewed medical records to confirm the diagnosis confirmation of SDR and TTN. All the criteria, including X-ray, laboratory assessments, clinical evolution, aimed to exclude other causes of ventilatory support as sepsis, meconium aspiration syndrome, birth with acidosis.

4

Expected result

We expect to identify SDR and TTN with a set of criteria based on scientific references.