VERSION 2

NOV 27, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocol.s.io.9tyh6pw

External link:
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2786677/>

Protocol Citation: Ian M Mackay, Judy A Northill 2019. Respiratory picornavirus genotyping conventional nested RT-PCR ("Wisdom VP42 assay"). protocols.io <https://dx.doi.org/10.17504/rotocols.io.9tyh6pw>

MANUSCRIPT CITATION:
Reference 1. Screening Respiratory Samples for Detection of Human Rhinoviruses (HRVs) and Enteroviruses: Comprehensive VP4-VP2 Typing Reveals High Incidence and Genetic Diversity of HRV Species C. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2786677/>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Nov 27, 2019

Last Modified: Nov 27, 2019

PROTOCOL integer ID:
30296

Keywords: enterovirus, rhinovirus, EV-D68, EV-A71, RV-C, RV-A, RV-B, picornavirus, respiratory, subgenomic, conventional RT-PCR, RT-PCR, nested PCR

Respiratory picornavirus genotyping conventional nested RT-PCR ("Wisdom VP42 assay") V.2

Version 1 is forked from [Respiratory picornavirus genotyping conventional nested RT-PCR \("Wisdom VP42 assay"\)](#)

Ian M Mackay¹, Judy A Northill¹

¹Public Health Virology, Forensic and Scientific Services

Ian M Mackay
Public Health Virology, Forensic and Scientific Services

ABSTRACT

This is my preferred, previously published [Ref 1], rhinovirus (RV) and enterovirus (EV) genotyping assay when seeking to identify the genotype of a respiratory picornavirus detected in a clinical sample extract. It is employed after use of a screening real-time RT-PCR has identified a respiratory picornavirus.

I have not confirmed that it can detect every single RV genotype but I do know that it detects *many* from each of the three RV species (*Human rhinovirus A*, *Human rhinovirus B* and *Human rhinovirus C*) as well as at least some *Human enterovirus* (EV) genotypes.

The assay picks up EVs due to the shared genetic similarities in the 5'UTR target region. EVs can be discriminated using subgenomic sequencing (see VP42 typing assay protocol), or simply described as 'respiratory EVs' since there is no specific-specific vaccine or treatment available anyway.

This is a robust primary subgenomic sequencing assay. It is more sensitive than any VP1 protocols because it targets more conserved primer target sites. It produces a more reliable typing result than does the 5'UTR region alone.

MATERIALS

MATERIALS

MyFi Mix **Bioline Catalog #BIO-25049**

SuperScript™ III One-Step RT-PCR System with Platinum™ Taq DNA Polymerase **Thermo Fisher Catalog #12574026**

STEP MATERIALS

SuperScript III One-Step RT-PCR System with Platinum Taq **Invitrogen - Thermo Fisher Catalog #12457-026**

MyFi Mix **Bioline Catalog #BIO-25049**

Oligonucleotide sequences...

1		Name	Sequence (5'-3')
	RT-PCR (Round 1)	HRV_HEV VP42 OS	CCGCCCCCTGAATGYGGCTAA
		HRV_HEV VP42 OAS	ACATRTTTYTNNCCAAANAYDCCCAT
	PCR (Round 1)	HRV_HEV VP42 IS	ACCRACACTTTGGGTGTCCGTG

	HRV_HEV VP42 IAS	TCWGGHARYTTCCAMCACCANCC
--	------------------	-------------------------

1. Expected amplicon sizes: Round 1: ~380 base pairs; Round 2:
2. The naming used here is my in-house adaptation (FYI: 01 - forward / sense; 02 - reverse / antisense; .x - version of the design of this particular named oligonucleotide). If you prefer to be true to the original publication, please see Ref 1

Reagents

2

☒ SuperScript III One-Step RT-PCR System with Platinum Taq **Invitrogen - Thermo Fisher Catalog #12457-026**

☒ MyFi Mix **Bioline Catalog #BIO-25049**

Reaction set-up

3

ROUND 1 REACTION MIX

- Ideally, set up more reaction mixes than you'll need for both rounds at the same time
- Strips of 8x 0.2ml tubes are good for this use
- Freeze in a frost-free freezer at -20°C until needed

Reagent	Vol (μl) 1x	Final reaction concentration
Nuclease-free water	4.08	N/A
Reaction Mix (2X)	10.00	1X
HRV_HEV VP42 OS 200pmol/μl	0.06	600nM
HRV_HEV VP42 OAS 200pmol/μl	0.06	600nM
SuperScript® III RT/ Platinum® Taq Enzyme Mix	0.20	Unknown
Template	5.00	N/A

1. Dispense 15μL to each reaction tube
2. Total reaction volume will be 20μl

Reagent	Vol (μl) 1x	Final reaction concentration
Nuclease-free water	7.92	N/A
MyTaq Reaction Buffer (2X)	10.00	1X
HRV_HEV VP42 IS 200pmol/μl	0.038	380nM
HRV_HEV VP42 IAS 200pmol/μl	0.038	4.75mM
1st round amplicon	2.00	N/A

1. Dispense 18μL into each reaction tube
2. Total reaction volume will be 20μl

Amplification: ROUND 1, RT-PCR

4

- This reaction has been run using a SimpliAmp (Applied Biosystems) thermal cycler
- Transfer 5μl of nucleic acid extract (extracted RNA, controls or NTC [nuclease-free water]) to defrosted reaction tubes and cycle

55°C	20 min	1X
94°C	2 min	1X
94°C	30 sec	
50°C	45 sec	4X
68°C	50 sec	
94°C	30 sec	
40°C	45 sec	25X
68°C	50 sec	

68°C	5 min	1X
15°C	∞	

Amplification: ROUND 2, PCR

- 5
- This reaction has been run using a SimpliAmp (Applied Biosystems) thermal cycler
 - Transfer 2µl of 1:100 pre-diluted Round 1 amplicon into defrosted 0.2ml reaction tubes and cycle

95°C	1 min	1X
95°C	30 sec	
50°C	45 sec	4X
72°C	50 sec	
95°C	30 sec	
40°C	45 sec	25X
72°C	50 sec	
72°C	5 min	1X
15°C	∞	

Amplicon visualisation

- 6
- electrophorese 2-5µl amplicon on a 1.5% agarose gel in 0.5X TBE buffer