

Oct 24, 2020

Restriction Digest

DOI

dx.doi.org/10.17504/protocols.io.bnwvmfe6

Zhujun Wei¹

¹2020 iGEM NEFU China

2020 iGEM NEFU China



Zhujun Wei

OPEN ACCESS



DOI: dx.doi.org/10.17504/protocols.io.bnwvmfe6

Protocol Citation: Zhujun Wei 2020. Restriction Digest. protocols.io https://dx.doi.org/10.17504/protocols.io.bnwvmfe6

License: This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 24, 2020

Last Modified: October 24, 2020

Protocol Integer ID: 43701

Abstract

This protocol includes three different methods of digestion.

Materials

Restriction enzyme Buffer 2.1/3.1/CutSmart



Safety warnings



• Please wear gloves for the experiment.

1 Add the following reagents to a tube.

Total		10ul	30ul	50ul
DNA		0.4ug	1.2ug	2ug
Enzyme		0.2ul	0.6ul	1ul
Buffer(10*)		1ul	3ul	5ul
ddwater Fill th water	e rest with			

- 2 Pipette up and down thoroughly.
- 3 Put at 37°C for about 2 hours (according to different enzymes).
- 4 Add the following reagents to a tube.

Total	10ul	30ul	50ul
DNA	0.4ug	1.2ug	2ug
Enzyme	0.2ul	0.6ul	1ul
Buffer(10*)	1ul	3ul	5ul
ddWater	Fill the rest with water	Fill the rest with water	Fill the rest with water

- 5 Pipette up and down thoroughly.
- 6 Put at 37°C for about 2 hours (according to different enzymes).
- 7 Put at 65°C for about 20 min.(according to different enzymes)
- 8 Purification of enzyme digestion products with Cycle Pure Kit D6492 (OMEGA bio-tek)



- 9 DNA product concentrations were measured using Nano
- 10 Add the following reagents to a tube:

Total	10ul	30ul	50ul
DNA product	0.4ug	1.2ug	2ug
CIP(NEB)	0.2ul	0.6ul	1ul
Cutsmart(10*)	1ul	3ul	5ul
ddWater	Fill the rest with water	Fill the rest with water	Fill the rest with water

- 11 Pipette up and down thoroughly.
- 12 Put at 37°C for about 2 hours (according to different enzymes).
- 13 Add the following reagents to a tube.

	Total	10ul	30ul	50ul
	DNA	0.4ug	1.2ug	2ug
	Enzyme 1	0.2ul	0.6ul	1ul
	Enzyme 2	0.2ul	0.6ul	1ul
Г	Buffer(10*)	1ul	3ul	5ul
	ddwater	Fill the rest with water	Fill the rest with water	Fill the rest with water

- 14 Pipette up and down thoroughly.
- 15 Put at 37°C for about 2 hours (according to different enzymes).



16 Add the following reagents to a tube.

Total	10ul	30ul	50ul
DNA product	0.4ug	1.2ug	2ug
DpnI(NEB)	0.2ul	0.6ul	1ul
Cutsmart(10*	1ul	3ul	5ul
ddwater	Fill the rest with water	Fill the rest with water	Fill the rest with water

- 17 Pipette up and down thoroughly.
- 18 Put at 37°C for about 2 hours (according to different enzymes).