OCT 16, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.8bjhskn

Protocol Citation: Igem Dusseldorf 2019. Reverse transcription of RNA to cDNA. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.8bjhskn>

MANUSCRIPT CITATION:
https://assets.thermofisher.com/TFS-Assets/LSG/manuals/MAN0012757_RevertAid_Reverse_Transcriptase_UG.pdf,
https://assets.thermofisher.com/TFS-Assets/LSG/manuals/MAN0012904_DyNAmo_ColorFlash_SYBR_Green_qPCR_F416L_UG.pdf

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 16, 2019

🌐 Reverse transcription of RNA to cDNA

Igem Dusseldorf¹

¹Heinrich-Heine Universität Düsseldorf

Igem Dusseldorf

ABSTRACT

Synthesizing cDNA (for qPCR)

GUIDELINES

Always work with gloves and safety gear and work on ice

MATERIALS

- 100 ng RNA
- 40x yellow sample buffer (from DyNAmo Color Flash Kit)
- random hexamer primers
- 5x RT-buffer
- dNTP Mix (10 mM of each nucleotide)
- RevertAid Reverse Transcriptase
- **You will require 100 ng for RT and 100 ng for -RT control (include a -RT control for each sample!)**

Preparation of 1.33X yellow buffer

- 1 33,25 μ L 40x yellow sample buffer + 966,75 μ L H₂O

optional: This was used to be able to see the pipetting scheme for qPCR more easily

Start

- 2 In PCR stripes, pipet in the following order:

100 ng template RNA
1 μ L random hexamer primers
RNase-free H₂O to 13 μ L volume

- 3 Prepare a Mastermix of the following reagents:

4 μ L 5x RT-buffer
2 μ L dNTP Mix (10 mM of each nucleotide)
1 μ L RevertAid Reverse Transcriptase

- 4 Add 7 μ L Mastermix to each reaction

- 5 For your -RT control, pipet 100 ng RNA in PCR stripes, leave out buffer and everything, just add H₂O to a final volume of 20 μ L. Mark as -RT so you can distinguish it from your actual cDNA!

PCR

- 6

PCR protocol:

10 min	25 °C
-----------	-------

60 min	42 °C
10 min	70 °C

7 Add 60 µL 1,33 x yellow sample buffer to each reaction

optional: This was used to be able to see the pipetting scheme for qPCR more easily

8 cDNA can be stored at 4 °C for a short time, otherwise, freeze at -20 °C