VERSION 3

OCT 15, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.78uhrww

Protocol Citation: Lutz Berwanger 2019. RNA extraction with PGTX.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.78uhrww>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 15, 2019

Last Modified: Oct 15, 2019

PROTOCOL integer ID:
 28660

RNA extraction with PGTX V.3

Lutz Berwanger¹

¹Institute of Synthetic Microbiology, Heinrich Heine University Düsseldorf

Axmann Lab

CyanoWorld

Lutz LCB Berwanger

Heinrich-Heine Universität Düsseldorf

ABSTRACT

This protocol is used for RNA extraction in cyanobacteria after Pinto et al. 2009.

Pinto, Fernando Lopes; Thapper, Anders; Sontheim, Wolfgang; Lindblad, Peter (2009): Analysis of current and alternative phenol based RNA extraction methodologies for cyanobacteria. In: *BMC molecular biology* 10, S. 79. DOI: 10.1186/1471-2199-10-79

MATERIALS

STEP MATERIALS

⊗ Ethyl alcohol, Pure 200 proof, for molecular biology **Merck MilliporeSigma (Sigma-Aldrich) Catalog #E7023**

⊗ Roti®-C/I **Carl Roth**

⊗ Isopropanol **Contributed by users**

⊗ PGTX **Contributed by users**

SAFETY WARNINGS

Additional Information

Always provide RNA work on ice and always wear Gloves.

GHS Label elements, including precautionary statements

Signal word Danger

Hazard statement(s)

H227 Combustible liquid.
H302 Harmful if swallowed.
H319 Causes serious eye irritation.
H311 Toxic in contact with skin.
H314 Causes severe skin burns and eye damage.
H330 Fatal if inhaled.
H341 Suspected of causing genetic defects.
H371 May cause damage to organs.
H373 May cause damage to organs through prolonged or repeated exposure.

H402 Harmful to aquatic life.

Precaution Statement(s)

P260 Do not breathe dust/ fume/ gas/ mist/ vapours/ spray.
P280 Wear protective gloves/ protective clothing/ eye protection/ face protection.
P284 Wear respiratory protection.
P305 + P351 + P338 IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P310 Immediately call a POISON CENTER or doctor/ physician.

Material list

Che mical	Volu me/ Mass	Com pany	Seria l no.	Com ment s
Phen ol	39.6g			See add. Info
Glycer ol	6.9m L			

8-hydroxyquinoline	0.1g			
EDTA	0.58g			See add. Info
Sodium acetate	0.8g			
Guanidine thiocyanate	9.5g			
Guanidine hydrochloride	4.6g			
Chloroform				See add. Info
Isopropanol				See add. Info
EtOH	70%			
DEPC treated Water		Roth		

Buffer

Name	Ingredients		Comments
100 mL PGTX Solution:	Phenol Glycerol 8-hydroxyquinoline EDTA Sodium acetate Guanidine thiocyanate Guanidine hydrochloride	39.6 g 6.9 mL 0.1 g 0.58 g 0.8 g 9.5 g 4.6 g	

Equipment list

Device	Company	Comments
Centrifuge	Eppendorf	4°C

Thermo Mix	Eppendorf	95°C
Centrifuge	Heraeus	4°C
Freezer	Liebherr	-20°C

- 1 Fill 50 mL Falcon tube with ice. Fill with bacterial liquid culture, up to a volume of ~45 mL.
- 2 Centrifuge for 3 minutes at 4°C and 4.000 g.
⌚ 00:03:00
- 3 Discard supernatant. Resuspend cell pellet in residual water (~1 mL). Transfer to 2 mL 'safe lock' tube.
- 4 ⌚ 00:00:15
- 5 Resuspend the cell pellet in 1 ml of PGTX. Freeze in liquid nitrogen and store at -20 °C.
⊗ PGTX Contributed by users

Safety information

Wear goggles, a lab coat and gloves when dealing with PGTX and liquid nitrogen.

- 6 Incubate for 5 min at 95 °C, shaking at 250 rpm in Thermomixer (Eppendorf)
⌚ 00:05:00
- 7 Rapidly chill 5 min on ice.

⌚ 00:05:00

8 Add 700 µl of Roti-C/I

⊗ Roti®-C/I **Carl Roth**

Safety information

Wear goggles, a lab coat and gloves when dealing with phenol and chloroform.

9 Let the samples incubate for 10 min at room temperature in a Thermomixer. Vortex from time to time.

⌚ 00:10:00

10 Centrifuge for 15 min at 14.000 g, 4 °C. Transfer the upper aqueous phase (**contains RNA**) to a fresh reaction tube and add the same volume (450 µL) of Aqua- C/I (Chloroform/Isoamylalcohol).

⌚ 00:15:00

Safety information

Wear goggles, a lab coat and gloves when dealing with phenol and chloroform.

11 Thoroughly mix by vortexing. Centrifuge for 15 min at high speed. Transfer the upper aqueous phase to a 1.5 mL reaction tube.

⌚ 00:15:00

Safety information

Wear goggles, a lab coat and gloves when dealing with phenol and chloroform.

12 Add 1 volume of isopropanol.

⊗ Isopropanol **Contributed by users**

13 Mix and incubate for at least 30 min at -20 °C. (Can be left overnight.)

⌚ 00:30:00

14 Centrifuge at 14.000 g for 30 min.

00:30:00

15 Discard supernatant.

16 Wash with 75% chilled ethanol. Avoid resuspending the pellet.

Ethyl alcohol, Pure 200 proof, for molecular biology **Merck MilliporeSigma (Sigma-Aldrich) Catalog #E7023**

17 Centrifuge at 14.000 g.

00:05:00

18 Repeat washing step with 75% chilled ethanol.

Note

Remove excess of EtOH by using a pipet.

19 Air dry pellet at RT. Do not overdry!

00:10:00

20 Resuspend the pellet with 30 μL volume of ddH₂O.

Note

Usually 40 μL DECP-treated H₂O