AUG 19, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.4vygw7w

Protocol Citation: Beijing Genomics Institute 2019. RNA Isolation from Plant Tissue Protocol 3: CTAB-PVP Method. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.4vygw7w>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jun 28, 2019

Last Modified: Aug 19, 2019

PROTOCOL integer ID:
 25240

Keywords: RNA, RNA isolation, RNA extraction, plant tissue, plant RNA

RNA Isolation from Plant Tissue Protocol 3: CTAB-PVP Method

In 1 collection

Beijing Genomics Institute¹

¹Beijing Genomics Institute

GigaScience Press

BGI

Eric Carpenter

ABSTRACT

Implemented by: Beijing Genomics Institute

Note

A similar protocol was used by C. dePamphilis and P. Ralph.

This protocol is part of a collection of eighteen protocols used to isolate total RNA from plant tissue. (RNA Isolation from Plant Tissue Collection: <https://www.protocols.io/view/rna-isolation-from-plant-tissue-439gyr6>)

ATTACHMENTS

[journal.pone.0050226.s01](#)
 1-1.pdf

MATERIALS

Reagents:

CTAB-PVP Buffer:

- CTAB (Hexadecyltrimethylammonium bromide; 2 % w/v)
- PVP-40 (2 % w/v)
- 100 mM Tris-HCl (pH 8.0)
- 25 mM EDTA
- 2 M NaCl (Warmed to 65 °C in a water bath to suspend in solution)
- Add β -ME to final concentration of 2 % before use

SSTE buffer:

- 1 M NaCl
- SDS (0.5 % w/v)
- 10 mM Tris-HCl (pH 8.0)
- EDTA (1 mM)

Other reagents:

- 75 % ethanol (treated with 0.1 % DEPC)
- 96–100 % ethanol
- Acid phenol (pH 4.5)
- Chloroform
- Isoamyl alcohol
- 10 M LiCl
- Glycogen (5 mg/ml)
- 3 M Sodium Acetate (pH 5.2)
- RNase free water

SAFETY WARNINGS

Please see SDS (Safety Data Sheet) for hazards and safety warnings.

- 1 Grind tissue to a powder in liquid nitrogen.
- 2 Add 200 mg – 500 mg of ground, frozen tissue to 3.0 mL of pre-heated [extraction buffer](#) in a 5 ml tube.

- 3 Vortex the tube until the tissue is mixed with the buffer.
- 4 Incubate the tube at 65 °C for 00:30:00 (min), vortexing briefly (00:00:15) every 2–3 min during the incubation.
- 5 Aliquot the mixture into four 2 ml RNase free tubes, 1 mL each tube.
- 6 Spin the tube at 12000 x g for 00:10:00 in a microcentrifuge.

Note

All of the insoluble matter should form a pellet at the bottom of the tube.

Note

When used for algae the centrifugation was performed at 3000 x g .

- 7 Pour the supernatant into a new 2 ml RNase free tube.

Note

When used for algae, 15 ml tubes were used.

- 8 Add equal volume of 24:1 chloroform:isoamyl alcohol to fill the tube.

9 Vortex tubes until the phases mix and appear cloudy.

9.1 Incubate at 20 °C – 25 °C for 00:05:00 .

10 Spin the tubes at 12000 x g for 00:10:00 in a microcentrifuge.

Note

When used for algae the centrifugation was performed at 3000 x g .

11 Transfer the upper, aqueous phase to new 2 ml RNase free tubes.

Note

When used for algae, 15 ml tubes were used.

11.1 Repeat step 7 to 9 one more time. go to step #7

12 Transfer the upper, aqueous phase to new 2 ml RNase free tubes.

12.1 Add 1/3 volume of 10 Molarity (M) LiCl to each tube.

12.2 Mix and let stand at 4 °C for 06:00:00 – 08:00:00 or overnight to precipitate RNA.

13 Spin the tubes at 18000 x g for 00:20:00 in a microcentrifuge.

13.1 Decant the supernatant, taking care not to lose the pellet.

14 Add 1 mL 75 % ethanol to the pellet.

15 Spin the tube at maximum speed (11270 x g) for 00:05:00 in a microcentrifuge.

15.1 Decant the supernatant carefully.

16 Repeat steps 14 and 15 one more time. go to step #14

17 Open cap and air-dry the pellet.

- 18** Add 30 µL RNase free water to dissolve the pellet.
- 18.1** Then add 70 µL [SSTE buffer](#) to each tube.
- 19** Combine all 4 tubes into 1 tube.
- 20** Add equal volume of 25:24:1 acid phenol:chloroform:isoamyl alcohol to the tube.
- 21** Vortex the tubes until the phases mix and appear cloudy.
- 21.1** Then incubate at 20 °C for 00:05:00 .
- 22** Spin the tube at 12000 x g for 00:10:00 .
- 23** Transfer the upper, aqueous phase to a new 2 ml RNase free tube.
- 23.1** Add equal volume of 24:1 chloroform:isoamyl alcohol to the tube.

24 Vortex the tubes until the phases mix and appear cloudy.

24.1 Then incubate at 20 °C for 00:05:00 .

25 Spin the tube at 12000 x g for 00:10:00 in a microcentrifuge.

26 Transfer the upper, aqueous phase to a new 2 ml RNase free tube.

26.1 Add 2 volumes of cooled 100 % ethanol, 1/10 volumes of NaAc (pH 5.2) and 2 µL glycogen.

26.2 Mix and incubate at -20 °C for 02:00:00 .

27 Spin the tube at 18000 x g for 00:20:00 in a microcentrifuge.

27.1 Then decant the supernatant, taking care not to lose the pellet.

28 Add 1 mL 75 % cooled ethanol to the pellet.

28.1 Leave at 20 °C for 00:03:00 .

29 Centrifuge at 4 °C for 00:05:00 at 12000 x g .

29.1 Decant the liquid carefully, taking care not to lose the pellet.

29.2 Briefly centrifuge to collect the residual liquid and remove it with a pipette.

30 Wash the pellet twice with cooled 75 % DEPC-ethanol.

30.1 Open cap and air-dry the pellet.

31 Add 30 µL RNase-free water to dissolve the pellet.

32 Treat RNA with DNase I as per supplier's protocols.