OCT 04, 2019

RPA from banana DNA

Théo Nass¹, Luc Gabel¹

¹EPFL - EPF Lausanne

iGEM EPFL

Théo Nass

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.5cdg2s6

Protocol Citation: Théo Nass, Luc Gabel 2019. RPA from banana DNA.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.5cdg2s6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
 We are still developing and optimizing this protocol

Created: Jul 10, 2019

Last Modified: Oct 04, 2019

PROTOCOL integer ID:
 25701

MATERIALS

- nuclease-free water/ double-distilled autoclaved water

TE buffer

- 10 mM Tris pH 8.0
- 1 mM EDTA

DNA extraction :

- Banana
- Dish soap
- 1g table salt
- 25 ml pre-chilled (4°C) 90% isopropyl alcohol (rubbing alcohol)
- 70% ethanol

Primers :

- Fwd : 5' CATTCTAATACGACTCACTATAGGGATTATCTGCAAAA AACTACGG 3'
- Rev : 5' TAAAAAGTGCTTCGGTGCAAAATAAGAAACGAT 3'

Rehydration solution

- 1.18 µl rehydration buffer
- 0.84 µl 10µM forward primer
- 0.84 µl 10µM reverse primer
- 1 µl 280 mM magnesium acetate
- 2 reaction pellets (included in the kit)
- DNA ladder
- Loading buffer
- 2% agarose gel

Banana DNA extraction

- 1 Put chopped banana in a closed ziploc-like bag

- 2 Add diluted 1:10 dish soap solution and 1g table salt to fruits
- 3 Mash it until homogeneous mixture is obtained
- 4 Strain the mixture through a household coffee filter
- 5 Add 25 ml isopropyl alcohol to strained liquid
- 6 Let it sit for 5 min until phase separation
- 7 Collect upper white layer (precipitated DNA) with a toothpick, place it on a new coffee filter and wash it with 70% ethanol
- 8 Pat it dry with paper towels
- 9 Dissolve DNA in 1 ml TE buffer

Nanodrop

- 10 Perform a NanoDrop analysis

Storage

- 11 Set 1 μl aside; store rest at 4°C

Prepare amplification mix

- 12 Prepare the rehydration solution for the RPA

	experiment	RPA control
Rehydration buffer	5.9 μl	5.9 μl
10 μM forward primer	0.42 μl	0.42 μl

10 μ M reverse primer	0.42 μ l	0.42 μ l
280 mM ammonium acetate	0.5 μ l	0.5 μ l
Sample extract	1 μ l	0 μ l
Water	1.76 μ l	2.76 μ l
Total	10 μ l	10 μ l

13 Vortex and spin briefly

14 Rehydrate 1 RPA reaction pellet in each one of the rehydration solutions

15 Pipette up and down to resuspend pellet

16 Vortex and spin briefly

RPA reaction

17 Heat the reaction at 39°C for 20 minutes.

DNA purification step

18 should do a standard DNA purif kit (which we don't have at the moment) so we'll try the same purif as we did for the banana DNA

Gel electrophoresis

19 Run a gel electrophoresis:

Lane	1	2	3
Sample	DNA ladder	Sample	PCR control