

Aug 13, 2020

Salmonella live attenuated vaccine preparation.

DOI

dx.doi.org/10.17504/protocols.io.bjpdkmi6

Angel A Justiz-Vaillant¹, Suzette Curtello²

¹University of the West Indies St. Augustine; ²University of the West Indies. Mona Campus.

University of the West In...

angel.vaillant@sta.uwi.e...



Angel A Justiz-Vaillant

University of the West Indies St. Augustine

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.bjpdkmi6>

Protocol Citation: Angel A Justiz-Vaillant, Suzette Curtello 2020. Salmonella live attenuated vaccine preparation.. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bjpdkmi6>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited



Protocol status: Working

We use this protocol and it's working

Created: August 13, 2020

Last Modified: August 13, 2020

Protocol Integer ID: 40389

Keywords: attenuated vaccine preparation, salmonella, vaccine preparation

Troubleshooting

- 1 A Salmonella cocktail (mixture of five Salmonella strains: S. Montevideo, S. Yeerongpilly, S. Augustenborg, S. Kentucky and S. Typhimurium) is incubated at 37°C in peptone broth with combinations of natural extracts.
- 2 Preparation includes clean, sun dried onion and garlic cloves.
- 3 After a week of drying, onion and the cloves of garlic is separately grounded using electric blender,
- 4 Dilute the plant preparations in 1L distilled water.
- 5 Each extract is used individually and in combined concentration.
- 6 Five ml of the diluted extract (15 mg/ml) is incorporated in differential and selective Salmonella plating media.
- 7 A bacteriogram is carried out to check for bacteria attenuation.
- 8 The immunogen contains 5 serovars of Salmonella (10^8 CFU of vaccine strain) per chicken in 1 ml of incomplete Freund's adjuvant.
- 9 It is given intramuscularly at chicken breast
- 10 The immunogen (5 serovars) without incomplete Freund's adjuvant (IFA) is given orally (10^8 CFU per chicken).
- 11 All Salmonella isolates were investigated for their antibiotic resistance with the disc diffusion test using the following discs (Difco):
- 12 gentamicin (10 µg), kanamycin (30 µg), ampicillin (10 µg), amikacin (30 µg), trimethoprim/sulfamethoxazole (1.25/23.75 µg), chloramphenicol (30 µg), cefazolin (30 µg), cephalothin (30 µg), cefepime (30 µg), cefotaxime (30 µg), streptomycin (10 µg), cefoxitin (30 µg), nalidixic acid (30 µg), ciprofloxacin (5 µg), norfloxacin (10 µg), tetracycline (30 µg) and imipenem (10 µg).



- 13 The five Salmonella serovars were fully attenuated and inoculated in appropriate media and store at -20 degree Celsius.