FEB 27, 2019

Sanger Sequencing

In 1 collection

Celeste M M.

Karch¹,

Rita Martinez¹, Jacob Marsh¹

¹Washington University in St Louis

Neurodegeneration Method Development Community

Tech. support email: ndcn-help@chanzuckerberg.com

Celeste M M. Karch

Washington University in St Louis

ATTACHMENTS

[Comprehensive Genomic
Editing and Screening
Protocol Updated
02142019.docx](#)

GUIDELINES

This protocols is part of the [Screening Edited iPSC Clones collection](#).

SAFETY WARNINGS

Please refer to the SDS (Safety Data Sheet) for information about hazards, and to obtain advice on safety precautions.

BEFORE START INSTRUCTIONS

After identifying iPSC clones with the appropriate banding pattern, gDNA from iPSC clones of interest should be sequenced in order to confirm genotype. To do this, set up a sequencing reaction as outlined below.

Perform PCR on gDNA using the following TouchDown Protocol (50µL reaction). The PCR amplicon size should be between 300-400 bp. The primers used for the RFLP screening can also be used for Sanger Sequencing.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.x8tfrwn

Protocol Citation: Celeste M M. Karch, Rita Martinez, Jacob Marsh 2019. Sanger Sequencing. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.x8tfrwn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 17, 2019

Last Modified: Feb 27, 2019

PROTOCOL integer ID:
20467

PCR

- 1 Set up PCR on ice, add reagents in desired order (however it is best to add the polymerase and

gDNA last).

	Volume	x# rxns
5x Green GoTaq Flexi Buffer	10 μ l	
25mM MgCl ₂	6 μ l	
25mM dNTPs	0.8 μ l	
Forward Primer (10 μ M)	2 μ l	
Reverse Primer (10 μ M)	2 μ l	
GoTaq DNA Polymerase (5U/ μ L)	0.25 μ l	
Milli-Q H ₂ O	26.95 μ l	
QuickExtract gDNA	2 μ l	
Total	50 μ l	

Segment	Cycle s	Temperature	Time
1	1	94°C	5 minutes
2	10	94°C	30 seconds
		65°C - 1°C/ cycle	30 seconds
		72°C	1 minute
3	35	94°C	30 seconds
		55°C	30 seconds
		72°C	1 minute
4	1	72°C	10 minutes
5	1	4°C	Forever

After product has been run in the thermocycler

- Run 15 μ L of each sample on a 2% gel to determine presence of PCR product.

3 Run product on gel at 150 V for 01:30:00 .

4 Image gel.

5 Save image.

6 After confirming the presence of PCR product, move on to sequencing reaction protocol below:

	Volume	x# rxns
PRC product	1 µl	
Primer (10µM)	1 µl	
Milli-Q H2O	10 µl	
Total	12 µl	

Send out a Forward and a Reverse for each sample.