


Aug 09, 2025

Version 5

SARS-CoV-2 nsp3 macrodomain Time-Resolved FRET peptide displacement assay V.5

DOI

<https://dx.doi.org/10.17504/protocols.io.eq2ly7r2mlx9/v5>



Haim Barr^{1,2}, Noa Lahav^{1,2}

¹The Weizmann Institute of Science; ²ASAP Discovery Consortium

Haim Barr: General acknowledgement: The Wohl Drug Discovery institute, The Nancy and Stephen Grand Israel National Center for Personalized Medicine.;

ASAP Discovery



Noa Lahav

Weizmann Institute

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.eq2ly7r2mlx9/v5>

Protocol Citation: Haim Barr, Noa Lahav 2025. SARS-CoV-2 nsp3 macrodomain Time-Resolved FRET peptide displacement assay. protocols.io <https://dx.doi.org/10.17504/protocols.io.eq2ly7r2mlx9/v5> Version created by **Nick Lynch**

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: June 08, 2025

Last Modified: August 09, 2025

Protocol Integer ID: 219785

Keywords: Coronaviridae, Time-Resolved FRET assay, Fluorescence assay, Mac1, Nsp3, TR-FRET, HTRF, ADPr, Displacement, Screening, Assay, Inhibitor, Fragment, Binding, Macrodomain, SARS-CoV-2, resolved fret peptide displacement assay, fret peptide displacement assay this protocol detail, peptide, modified peptide, binding of adenosine, affinity of mac1, assay for sar, sars cov2 mac1183000, potential inhibitor, adenosine, inhibitor, sar

Funders Acknowledgements:

National Institutes of Health/National Institute Of Allergy and Infectious Diseases (NIH/NIAID)

Grant ID: U19AI171399

Disclaimer

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Abstract

This protocol details the Time-Resolved FRET (TR-FRET) assay for SARS-CoV-2 nsp3 (*non-structural protein 3*) macrodomain (Mac1) binding of adenosine diphosphate (ADP)-ribosylated (ADPr) peptide. This method measures the affinity of Mac1 to its potential inhibitors by using a specific ADPr-modified peptide that allows the detection of binding. When bound, the biotinylated-peptide and the HIS-tagged Mac1 form a proximity complex that is detected by TR-FRET using Streptavidin-Eu Cryptate and anti-HIS-XL665 as a donor/acceptor pair. Excitation of the Eu Cryptate complex at 325 nm emits a resonant energy of 625 nm which in turn excites the XL665 to emit fluorescence at 665 nm. This energy transfer occurs only when ADPr modified peptide is in sufficient proximity to Mac1 while inhibitors which displace the peptide will prevent energy transfer. Binding is reported as the ratio of Acceptor/Donor (E_m/E_m) X 10,000.

Experiment Concentrations (From Stock to Assay)

	A	B	C	D	E
	Reagent	Stock	Loaded into Combi	Final in assay plate	Units
	His-SARS CoV-2 MAC1	183000	50	12.5	nM
	Substrate (Biotin-ADPr)	10000000	1600	400	nM
	Detection solution				
	Streptavidin-XL665 (SA-XL)	1	0.25	0.125	%
	MAb Anti-6HIS-Eu cryptate Gold	100	0.25	0.125	%
	Assay buffer				
	HEPES pH=7.0	250	25	25	mM
	NaCl	200	20	20	mM
	BSA	0.5	0.05	0.05	mg/ml
	Tween 20	0.5	0.05	0.05	%
	HTRF PPI Europium Detection Buffer	100	10	10	%

For more information, please check out the "Materials" Section

Compound Plate Design for Dose Response:

Total Assay Volume: 16 μ L

Compounds Top Assay Concentration: 100 μ M

Dilution Factor: 3

Dose Response Points: 10

Number of Replicates: 2

Backfill with DMSO: Yes

Compounds Plate Design for 2-Point Assay:

Total Assay Volume: 16 μ L

Compounds Assay Concentration: 100 μ M and 50 μ M

Dilution Factor: 2

Dose Response Points: 2

Number of Replicates: 2

Backfill with DMSO: Yes



Materials

Assay Buffer Reagents (Concentration listed are from Stock Solutions)

1. [M] 250 millimolar (mM) ☒ HEPES 0.5M buffer soln. pH 7.0 Fisher Scientific Catalog #AAJ60064AE (or similar)
2. [M] 200 millimolar (mM) ☒ Sodium chloride Merck MilliporeSigma (Sigma-Aldrich) Catalog #S9888 (or similar)
3. [M] 0.5 mg/mL ☒ Bovine Serum Albumin (BSA) Merck MilliporeSigma (Sigma-Aldrich) Catalog #A7030
4. [M] 0.5 % volume ☒ TWEEN® 20 Merck MilliporeSigma (Sigma-Aldrich) Catalog #P9416
5. [M] 100 % volume ☒ HTRF PPI Europium Detection Buffer CISBIO BIOASSAYS (PerkinElmer) Catalog #61DB9RDF

***Note:** There are several forms of the Assay Buffer in this experiment. The **Assay Buffer** is the final, active buffer used throughout the experiment and has all of the five above reagents included. **HTRF PPI Europium Detection Buffer** needs to be added fresh before each experiment. Thus, there was an intermediate Buffer called **Mac1 Buffer** that contained HEPES, NaCl, BSA, and Tween only. Mac1 Buffer was filtered and stored at 4°C. **HTRF PPI buffer** was then added to Mac1 Buffer fresh (to a final concentration of 10%) prior to performing the experiment--creating the **active Assay Buffer**.

Detection Solution Reagents (Concentration listed are from Stock Solutions)

[M] 1 % volume ☒ Streptavidin-XL665 CISBIO BIOASSAYS (PerkinElmer) Catalog #610SAXAC

- **Note:** Streptavidin-XL665 was dissolved in triply distilled water and diluted with HTRF PPI buffer to its stock concentration and then was aliquoted into 1.5mL sterile conical tubes

[M] 100 Mass Percent

☒ MAb Anti-6HIS-Eu cryptate Gold CISBIO BIOASSAYS (PerkinElmer) Catalog #61HI2KLA

Note: MAb Anti-6HIS-Eu cryptate Gold was dissolved in triple distilled water and then aliquoted into 1.5mL sterile conical tubes

Additional Reagents:

[M] 183000 nanomolar (nM) **His-SARS COV2 MAC1 Enzyme**

- The Enzyme stock was originally [M] 183000 nanomolar (nM) and was diluted to [M] 50 nanomolar (nM) before every experiment in **freshly made Assay Buffer**. The final assay concentration is

[M] 12.5 nanomolar (nM)

[M] 10000000 nanomolar (nM) **Substrate (Biotin-ADPr) MAC1**

- Substrate stock (ARTK(Bio)QTARK(Aoa-RADP)S) was dissolved in DMSO to the stock concentration. Before each experiment, the substrate stock was diluted to [M] 1600 nanomolar (nM) in freshly made Assay Buffer.

Protocol materials

 Sars Cov 2 Mac domain Plasmid **addgene Catalog #204787**

Safety warnings

 Please be sure to wear proper Personal Protective Equipment (PPE) while performing this experiment.

Before start

Note: Inhibitor compounds stock concentration is **[M] 20 millimolar (mM)** . Compounds are pre-dispensed into 384 plates and stored at -20°C until use.



SARS-CoV-2 nsp3 expression and purification

1 The protein used in this assay was expressed and purified using the following protocol.

We used the following plasmid in the protein expression and purification protocol.

 Sars Cov 2 Mac domain Plasmid **addgene Catalog #204787**

Protocol

NAME

SARS-CoV-2 nsp3 macrodomain His-tagged expression and purification protocol for assay

CREATED BY

korvus.wang

Preview

Prepare Reagents

2 **PREPARE** all of the reagents/buffers required for this experiment.

Assay Buffer

	A	B	C	D	E
	Reagent	Stock	Loaded into Combi	Final in assay plate	Units
	HEPES pH=7.0	250	25	25	mM
	NaCl	200	20	20	mM
	BSA	0.5	0.05	0.05	mg/ml
	Tween 20	0.5	0.05	0.05	%
	HTRF PPI Europium detection buffer	100	10	10	%

Reagents (dilute reagents in assay buffer for required volume)



	A	B	C	D	E
	Reagent	Stock	Loaded into Combi	Final in assay plate	Units
	His-SARS-CoV-2 MAC1	183000	50	12.5	nM
	Substrate (Biotin-ADPr)	10000000	1600	400	nM

Detection Solution (dilute reagents in assay buffer for required volume)

	A	B	C	D	E
	Reagent	Stock	Loaded into Combi	Final in assay plate	Units
	Streptavidin-XL665 (SA-XL)	1	0.25	0.125	%
	MAb Anti-6HIS-Eu cryptate Gold	100	0.25	0.125	%

Prepare 384-well Plate

16m

- 3 **PRIME** Multi-Drop Combi Tube Dispensing Cassette with **Assay Buffer** by selecting the **PRIME** button on the Combi Dispenser until the tubes are filled completely.
 - **Note:** Be sure to cycle dispensing several times on a clean plate lid (This confirms there are no bubbles in the Dispensing Cassette).
- 3.1 **DISPENSE**  4 µL Mac1 Buffer to Columns **1 and 23** of assay plate
 - **Note:** These will represent the ***inhibitor control columns***
- 3.2 **EMPTY** Multi-Drop Combi Tube Dispensing Cassette (by selecting the **EMPTY** button on the Combi Dispenser until the tubes of the cassette are emptied). **Discard** the **Assay Buffer** discharged from the cassette.
- 4 **PRIME** Multi-Drop Combi Tube Dispensing Cassette with **His-SARS COV2 MAC1 Enzyme** by selecting the **PRIME** button on the Combi Dispenser until the tubes are filled completely.
 - **Note:** Be sure to cycle dispensing several times on a clean plate lid (This confirms there are no bubbles in the Dispensing Cassette).



- 4.1 **DISPENSE**  4 μ L [M] 50 nanomolar (nM) His-SARS COV2 MAC1 Enzyme to Columns **2-22** and **24** of assay plate
- Note:**
- [M] 50 nanomolar (nM) His-SARS COV2 MAC1 is four times the final concentration for the assay. It will be diluted to be a final concentration of [M] 12.5 nanomolar (nM) His-SARS COV2 MAC1 Enzyme
 - Column **2** and Column **24** are **neutral control columns** (Contain: Enzyme, Substrate, DMSO; **no experimental compounds**)
- 4.2 **EMPTY** Multi-Drop Combi Tube Dispensing Cassette (by selecting the **EMPTY** button on the Combi Dispenser until the tubes of the cassette are emptied). **Discard** the [M] 50 nanomolar (nM) His-SARS COV2 MAC1 Enzyme discharged from the cassette.
- 5 **PRIME** Multi-Drop Combi Tube Dispensing Cassette with **Assay Buffer** by selecting the **PRIME** button on the Combi Dispenser until the tubes are filled completely. Then, **EMPTY** the Multi-Drop Combi Tube Dispensing Cassette (by selecting the **EMPTY** button on the Combi Dispenser until the tubes of the cassette are emptied). **Discard the Assay Buffer discharged from the cassette.**
- 6 **CENTRIFUGE** plate  1500 rpm, Room temperature, 00:01:00 to remove bubbles 1m
- 7 **INCUBATE** plate for  00:15:00 at  Room temperature 15m
- 8 **PRIME** Multi-Drop Combi Tube Dispensing Cassette with [M] 1600 nanomolar (nM) MAC1 Substrate (Biotin-ADPr) by selecting the **PRIME** button on the Combi Dispenser until the tubes are filled completely.
- **Note:** Be sure to cycle dispensing several times on a clean plate lid (This confirms there are no bubbles in the Dispensing Cassette).
- 8.1 **DISPENSE**  4 μ L [M] 1600 nanomolar (nM) MAC1 Substrate (Biotin-ADPr) into Columns **1-24** (full plate)
- Note:**
- [M] 1600 nanomolar (nM) MAC1 Substrate (Biotin-ADPr) is four times the final concentration for the assay. It will be diluted to be a final concentration of [M] 400 nanomolar (nM) MAC1 Substrate (Biotin-ADPr)



- 8.2 **EMPTY** Multi-Drop Combi Tube Dispensing Cassette (by selecting the **EMPTY** button on the Combi Dispenser until the tubes of the cassette are emptied). **Discard** the [M] 1600 nanomolar (nM) MAC1 Substrate (Biotin-ADPr) discharged from the cassette.
- 9 **CENTRIFUGE** plate 1500 rpm, Room temperature, 00:01:00 to remove bubbles
- 10 **PRIME** Multi-Drop Combi Tube Dispensing Cassette with **Assay Buffer** by selecting the **PRIME** button on the Combi Dispenser until the tubes are filled completely. Then, **EMPTY** the Multi-Drop Combi Tube Dispensing Cassette (by selecting the **EMPTY** button on the Combi Dispenser until the tubes of the cassette are emptied). **Discard the Assay Buffer discharged from the cassette.**
- 11 **PRIME** Multi-Drop Combi Tube Dispensing Cassette with [M] 0.25 % volume Detection Solution by selecting the **PRIME** button on the Combi Dispenser until the tubes are filled completely.
- **Note:** Be sure to cycle dispensing several times on a clean plate lid (This confirms there are no bubbles in the Dispensing Cassette).
- 11.1 **DISPENSE** 8 µL [M] 0.25 % volume Detection Solution into full plate
- Note:**
- [M] 0.25 % volume Detection Solution is two times the final concentration for the assay. It will be diluted in the plate to be a final concentration of [M] 0.125 % volume Detection Solution
- 11.2 **EMPTY** Multi-Drop Combi Tube Dispensing Cassette (by selecting the **EMPTY** button on the Combi Dispenser until the tubes of the cassette are emptied). **Discard** the [M] 1600 nanomolar (nM) MAC1 Substrate (Biotin-ADPr) discharged from the cassette.
- 12 **CENTRIFUGE** 1500 rpm, Room temperature, 00:01:00 plate to remove bubbles
- 13 **INCUBATE** plate for 01:00:00 at Room temperature
- Recommended:** Clean/Empty the Multi-Drop Combi Reagent Dispenser and Dispensing Cassette during this incubation step

1h



Read Plate Fluorescence

- 14 **READ** and **RECORD** the plate Relative fluorescence units (RFU) via the "**Mac1 Protocol**" on the PHERAstar FS Control Software.

Equipment

PHERAstar FS

Microplate reader

BMG LABTECH

0471B0001A

[https://www.bmglabtech.com/en/pherastar-fsx/?](https://www.bmglabtech.com/en/pherastar-fsx/?utm_term=pherastar%20plate%20reader&utm_campaign=usa.roi.products&utm_source=FwOoFR_5EUHUaAlkREALw_wcB)


[utm_term=pherastar%20plate%20reader&utm_campaign=usa.roi.products&utm_source=FwOoFR_5EUHUaAlkREALw_wcB](https://www.bmglabtech.com/en/pherastar-fsx/?utm_term=pherastar%20plate%20reader&utm_campaign=usa.roi.products&utm_source=FwOoFR_5EUHUaAlkREALw_wcB)

Expected result

Donor 325/620 ex/em should be ~ 5000 . Acceptor ~ 3000

Diagram of assay

- 15 Figure 1 graphical depiction of assay principal and its use in screening campaign




Principal of ADPr peptide displacement assay

A - Binding of ADPr-modified peptide to Mac1 protein **B**- Detection reagents added to protein+peptide complex **C**- TR-FRET detects binding of peptide to Mac1 if proximity of donor and acceptor detection reagents is sufficient to enable resonant energy transfer **D**- Inhibitor compounds are detected by reduced TR-FRET signal when inhibitor displaces/prevents binding of ADPr-peptide to Mac1

Experimental Design

16



384 plate layout

Protocol references

<https://doi.org/10.1126/sciadv.abf8711>