FEB 15, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.x59fq96

Protocol Citation: Alex Rajewski 2019. Seed Sterilization. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.x59fq96>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Feb 14, 2019

Last Modified: Feb 15, 2019

PROTOCOL integer ID:
 20385

Seed Sterilization

Alex Rajewski¹

¹University of California, Riverside

Alex Rajewski
 University of California, Riverside

ABSTRACT

This protocol details how to surface sterilize plant seeds for tissue culture regeneration/transformation or for other techniques requiring sterile seeds.

GUIDELINES

The ratios of bleach and HCl used here are very approximate and also determined empirically. If you wish, you can calculate the volume of HCl necessary to fill the chamber with 6.1% chlorine gas (according to doi:10.3791/56587)

MATERIALS

MATERIALS

- Petri Dish P212121 Catalog #LI-PD01100
- Hydrochloric Acid Contributed by users
- Sodium Hypochlorite Solution Contributed by users

- You will also need a chamber to contain the gas for the sterilization. We find that an old dessication chamber works well. Note the exposed metal will rust in this environment, so it is best to use an older chamber you don't care about.
- None of the vendors for the materials are important.
- The sodium hypochlorite solution (bleach) will vary in concentration between 5 and 9% depending on the vendor. We do not find that the protocol is sensitive to these variations, so use, whichever you have on hand.

SAFETY WARNINGS

- This protocol will create a chamber filled with chlorine gas, which is very toxic. It will also use concentrated HCl. Make sure that you have a proper fume hood to dispose of the gas and proper equipment to handle the HCl.

- 1 In a fume hood, place an empty dessication chamber with a petri dish of the seeds you wish to surface sterilize. Leave the lid of the petri dish halfway on, so that you can quickly close it later.

Note

- A dessication chamber per se is not necessary; you simply need some container to hold the gas for a few hours. It does not need to be thoroughly airtight, but it helps.
- If you plan to sterilize multiple genotypes of seeds, be aware that the gas produced will erase most markers. Pencil and paper labels work well though.

- 2 Place a 100mL beaker containing approximately 50 mL bleach in the chamber next to the petri dish of seeds.

3 Prepare approximately 3 mL HCl in a transfer pipet or similar.

4 The following steps must occur quickly to prevent the escape of the gas. Hold the lid to the chamber in one hand. With the other hand squirt the HCl into the beaker of bleach and quickly close the chamber to seal in the gas.

Note

The HCl and the bleach will react violently to produce the gas, do not be startled. The larger beaker should prevent any bubbles of bleach or HCl from contacting the rest of the set up.

5 Allow the seeds to be sterilized for approximately 3 hours. This time works for tomato, tobacco, and Arabidopsis seeds. The seeds may become whiter in color, this is also normal and does not seem to affect germination rates.

6 Once the time is up, quickly open the chamber and slide the lid completely onto the petri dish to seal it. The seeds can be stored like this for several weeks or transferred to a laminar flow hood for further use. Allow the chamber to air out for an hour or so.

7 If the reaction of the bleach and HCl went to perfect completion, the beaker should only contain salt water. In reality it is likely acidic. Dispose of this according to your local regulations.