JUN 04, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bg6kjzcv

Protocol Citation: Gabriele Kaminski Schierle 2020. Selecting the correct transfer membrane for Western blot enhances detection of alpha-synuclein and tau proteins. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bg6kjzcv>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jun 04, 2020

Last Modified: Jun 04, 2020

PROTOCOL integer ID:
 37804

Keywords: Western blot, gel transfer, alpha-synuclein, tau, PVDF, nitrocellulose, fixation

🌐 Selecting the correct transfer membrane for Western blot enhances detection of alpha-synuclein and tau proteins

📁 In 1 collection

Gabriele Kaminski Schierle¹

¹University of Cambridge

Molecular Neuroscience Group
 Tech. support email: gsk20@cam.ac.uk

Gabriele Kaminski Schierle
 University of Cambridge

ABSTRACT

The ability to detect proteins by Western blot, particularly from tissue samples, can be hampered by the choice of membrane and whether a fixation step is used. Here we compare three transfer membranes, polyvinylidene difluoride (PVDF) 0.2 μm, PVDF 0.45 μm and nitrocellulose 0.45 μm for the detection of rat α-synuclein from tissue and recombinant human tau. For α-synuclein fixation of the membrane directly after transfer was imperative for detection of the protein and use of the PVDF 0.45 μm membrane gave the highest detection signal. For tau, the signal was highest on nitrocellulose 0.45 μm and fixation did not enhance the signal.

MATERIALS

STEP MATERIALS

NuPAGE™ LDS Sample Buffer (4X) Thermo Fisher Catalog #NP0008
NuPAGE™ 4-12% Bis-Tris Protein Gels, 1.5 mm, 15-well Thermo Fisher Catalog #NP0336PK2
NuPAGE™ MES SDS Running Buffer (20X) Thermo Fisher Catalog #NP0002
Immobilon-PSQ PVDF Membrane 0.2um roll Merck MilliporeSigma (Sigma-Aldrich) Catalog #ISEQ00005
Immobilon-P PVDF Membrane, 0.45um, roll Merck MilliporeSigma (Sigma-Aldrich) Catalog #IPVH00010
Nitrocellulose Transfer Membrane Amersham plc Catalog #10600002
Methanol Contributed by users
NuPAGE™ transfer buffer Thermo Fisher Scientific Catalog #NP0006
PBS Contributed by users
Tween-20 Merck MilliporeSigma (Sigma-Aldrich) Catalog #P9416
Rabbit IgG horse radish peroxidase (HRP) linked Whole Ab Contributed by users Catalog #GENA934-1ML
SuperSignal™ West Pico PLUS Chemiluminescent Substrate Thermo Fisher Catalog #34579
Polyclonal Rabbit Anti-Human Tau Unconjugated Ig fraction Agilent Technologies Catalog #A002401-2
α -Synuclein (D37A6) XP® Rabbit mAb Cell Signaling Technology Catalog #4179
Rabbit IgG horse radish peroxidase (HRP) linked Whole Ab Contributed by users Catalog #GENA934-1ML

SAFETY WARNINGS

Follow safety guidelines and wear correct PPE when handling fixation solution containing 4% paraformaldehyde and 0.1% glutaraldehyde

1 Protein samples were boiled for 00:03:00 in 1x

NuPAGE™ LDS Sample Buffer (4X) **Thermo Fisher Catalog #NP0008**

and ran on

NuPAGE™ 4-12% Bis-Tris Protein Gels, 1.5 mm, 15-well **Thermo Fisher Catalog #NP0336PK2**

in

NuPAGE™ MES SDS Running Buffer (20X) **Thermo Fisher Catalog #NP0002**

using the

Equipment	
XCell SureLock Mini-Cell Electrophoresis System	NAME
Electrophoresis System	TYPE
Invitrogen	BRAND
EI0001	SKU

at 200 V for 00:30:00 .

- 2 Three membranes were selected to test Western blot transfer and detection, PVDF 0.2 μ m

Immobilon-PSQ PVDF Membrane 0.2um roll **Merck MilliporeSigma (Sigma-Aldrich) Catalog #ISEQ00005**

PVDF 0.45 μ m

Immobilon-P PVDF Membrane, 0.45um, roll **Merck MilliporeSigma (Sigma-Aldrich) Catalog #IPVH00010**

and nitrocellulose 0.45 μ m

Nitrocellulose Transfer Membrane **Amersham plc Catalog #10600002**

- 3 PVDF membranes were first activated with methanol by incubating for 00:03:00 .
Nitrocellulose membranes do not need to be activated first.

- 4 The gel was transferred onto each membrane using the

Equipment	
Invitrogen XCell II™ Blot Module	NAME
Western blot transfer module	TYPE
Invitrogen	BRAND
EI9051	SKU

with

NuPAGE™ transfer buffer **Thermo Fisher Scientific Catalog #NP0006**

+ 20%

Methanol **Contributed by users**

at 30 V for 01:30:00 . The blot module was kept On ice and surrounded by ice to keep cool and prevent protein damage.

5

Safety information

PFA and glutaraldehyde are toxic and should be handled in a chemical hood with appropriate PPE. The fixation solution should never be disposed of down the sink, but disposed of in a separate container for disposal through correct waste systems.

For detecting α -synuclein in tissue samples the membranes were fixed immediately after transfer with 4% paraformaldehyde (PFA), 0.1% glutaraldehyde in PBS for 00:30:00 . Lack of fixation leads to poor/no α -synuclein detection and fixation with only 4% PFA also leads to lower detection than with the addition of 0.1% glutaraldehyde.

6

Following disposal of the fixation solution into properly designated containers for removal, the membranes were blocked for 01:00:00 in 5% BSA in

PBS **Contributed by users**

+ 0.05%

Tween-20 **Merck MilliporeSigma (Sigma-Aldrich) Catalog #P9416**

(PBS-T) at Room temperature .

7

The membranes were then incubated either overnight or for 01:00:00 at

Room temperature in primary antibody probing for rat α -synuclein 1:1000 dilution

α -Synuclein (D37A6) XP® Rabbit mAb **Cell Signaling Technology Catalog #4179**

.

8

The membranes were washed three times for 00:05:00 in PBS-T.

9

The membranes were incubated for 01:00:00 at Room temperature in secondary antibody 1:4000

Rabbit IgG horse radish peroxidase (HRP) linked Whole Ab **Contributed by users Catalog**
#GENA934-1ML

10 The membranes were washed five times for 00:05:00 in PBS-T.

11 The membranes were developed in

SuperSignal™ West Pico PLUS Chemiluminescent Substrate **Thermo Fisher Catalog**
#34579

and imaged in a

Equipment

G:Box Chemi

NAME

Imaging system

TYPE

Syngene

BRAND

XX6

SKU

12

Expected result

The best signal for detection of alpha-synuclein from rat tissue was obtained on PVDF 0.45 μ m with fixation in 4% PFA and 0.1% glutaraldehyde.

- 13 For recombinant human tau, fixation was not required and the membranes were instead immediately blocked for 01:00:00 in 5% BSA in PBS-T at Room temperature .

- 14 The membranes were then incubated either overnight or for 01:00:00 at Room temperature in primary antibody probing for human tau 1:200 dilution

Polyclonal Rabbit Anti-Human Tau Unconjugated Ig fraction **Agilent Technologies Catalog**
#A002401-2

- 15

The membranes were washed three times for 00:05:00 in PBS-T.
- 16

The membranes were incubated for 01:00:00 at Room temperature in secondary antibody, 1:4000 dilution

Rabbit IgG horse radish peroxidase (HRP) linked Whole Ab **Contributed by users Catalog**
#GENA934-1ML
- 17

The membranes were washed five times for 00:05:00 in PBS-T.
- 18

The membranes were developed in Supersignal West Pico (#34580, ThermoFisher) and imaged in a

Equipment	
G:Box Chemi	NAME
Imaging system	TYPE
Syngene	BRAND
XX6	SKU

19

Expected result

The best signal for recombinant Tau was obtained on nitrocellulose 0.45 µm.

- 20 This optimisation was originally performed by Dr Colin Hockings, a former postdoc in the Molecular Neuroscience Group.