VERSION 1

SEP 17, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.7e5hjg6

Protocol Citation: Long Cai, Nina Dar, Yiing Lin, Shin Lin 2019. seqFISH Tissue Preservation. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.7e5hjg6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
 We are still developing and optimizing this protocol

Created: Sep 17, 2019

Last Modified: Sep 17, 2019

PROTOCOL integer ID:
 27837

Keywords: seqFISH, HUBMAP

seqFISH Tissue Preservation V.1

Long Cai¹, Nina Dar¹, Yiing Lin², Shin Lin³

¹California Institute of Technology; ²Washington University, Saint Louis; ³University of Washington

Human BioMolecular Atlas Program (HuBMAP) Method Development Community

Tech. support email: Jeff.spraggins@vanderbilt.edu

Shin Lin

ABSTRACT

Tissue Preparation for seqFISH

MATERIALS

Fixation

1 Clean surface with RNAzap. Make the following solutions.

1L PBS 1x, RNase-free

1. take 1 L RNase-free water bottle
2. use serologic pipet and draw up 100 mL water into another container
3. add 100 mL of 10x RNase-free PBS
4. mix thoroughly

50% sucrose

1. tare 1 L RNase-free disposable container
2. take 500 g RNase-free sucrose container
3. add RNase-free water to sucrose and transfer to 1 L container
4. continue to add RNase-free water until mass to 1 kg
5. mix thoroughly, may need agitation on platform at RT for some time
6. store at 4°C

NOTE: for fixation, minimum PFA to tissue volume is 20:1, ideal closer to 50:1

2 Prepare 4% PFA in 1x PBS, 120 mL total for 4 tissues 30 mL each in 50 mL conicals.

prepa re 3x conic als for 4 tissue s	
16% PFA	10 ml
10x PBS	4 ml
RNAs e free water	26 ml
total	40 ml

3 Cut freshly dissected tissue to small pieces for heart, try cutting slab, 2 cm x 1 cm x 0.4 cm thickness: 2 cm dimension from epicardium to endocardium.

- 4 Place each piece in a 50 ml tube (or smaller if you can cut to smaller pieces) in 4% PFA leave at RT for 3 hrs.
** the sample should be completely covered.
- 5 Wash three times with equal volume (30mL) of 1x PBS to remove PFA leave a bit of extra liquid in the bottom between each wash.

Sucrose gradient

- 6 Place the tissue in 10% Sucrose at RT and wait for it to sink in 50 mL conical, add 10 mL of sucrose wait for sample to sink to bottom – minimum incubation time of 30 min remove supernatant for next bath.
- 7 Place the tissue in 20% Sucrose at RT and wait for it to sink.
- 8 Place the tissue in 30% sucrose in 4°C – ready for shipping. Make sure there is enough 30% sucrose solution in container e.g. if in 50 mL conicals, have 50 mL of 30% sucrose.