



Jan 27, 2022

Version 4

Simple electroporation for efficient CRISPR/Cas9 genome editing in murine zygotes V.4

PLOS ONE

DOI

dx.doi.org/10.17504/protocols.io.b4dyqs7w

Simon E. Tröder¹, Branko Zevnik¹

¹CECAD, University of Cologne, Cologne, Germany



Simon E Tröder

Universität Köln



Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.b4dyqs7w>

External link: <https://doi.org/10.1371/journal.pone.0196891>

Protocol Citation: Simon E. Tröder, Branko Zevnik 2022. Simple electroporation for efficient CRISPR/Cas9 genome editing in murine zygotes. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.b4dyqs7w> Version created by **Simon E Tröder**

**Manuscript citation:**

Tröder SE, Ebert LK, Butt L, Assenmacher S, Schermer B, Zevnik B (2018) An optimized electroporation approach for efficient CRISPR/Cas9 genome editing in murine zygotes. PLoS ONE 13(5): e0196891. doi: 10.1371/journal.pone.0196891

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: January 27, 2022

Last Modified: January 27, 2022

Protocol Integer ID: 57496

Keywords: Zygote, Electroporation, CRISPR, Cas9, Cas12a, Mouse model, cas9 genome editing in murine zygotes electroporation, simple electroporation for efficient crispr, common electroporator with synthetic crispr, efficient crispr, genome editing in mice, cas9 genome editing, murine zygotes electroporation, mediated genome editing, other crispr nuclease, mice by electroporation, compatible with other crispr nuclease, synthetic crispr, crispr, current protocols for electroporation, simple electroporation, specialized electroporator, investment in specialized electroporator, electroporation, efficient introduction of specific mutation, common electroporator, rapid alternative to the elaborate pronuclear injection procedure, embryo viability, elaborate pronuclear injection procedure, genome, specific mutation, intact zygote, cas9 component

Abstract

Electroporation of zygotes represents a rapid alternative to the elaborate pronuclear injection procedure for CRISPR/Cas9-mediated genome editing in mice. However, current protocols for electroporation either require the investment in specialized electroporators or corrosive pre-treatment of zygotes which compromises embryo viability. Here, we describe an easily adaptable approach for highly efficient introduction of specific mutations in mice by electroporation of intact zygotes using a common electroporator with synthetic CRISPR/Cas9 components and minimal technical requirement. This protocol works efficiently with zygotes from a variety of genetic backgrounds and is compatible with other CRISPR nucleases like Cas12a.

Guidelines

Use only embryo-grade reagents.

Embryo culture media (M2, M16, KSOM/AA) may be purchased or prepared in-house as published (Behringer et al., 2014). Mouse pre-implantation embryos are incubated in pre-equilibrated M16 or KSOM/AA in a CO₂ incubator (5% CO₂, 37°C, 95% humidity).

Embryo-grade standard T₁₀E_{0.1} (10 mM Tris-HCl, 0.1 mM EDTA, pH 7.4) injection buffer is prepared as described (e.g., Chu et al., 2016).

Mouse zygotes are collected from the oviducts of superovulated females as described in published protocols (Behringer et al., 2014). To avoid developmental impairment is recommended to use only zygotes of high quality (i.e., presence of polar bodies and pronuclei).

This protocol works efficiently with zygotes from any source (generated by natural mating or IVF as well as freshly harvested or frozen-thawed) and from various genetic backgrounds like C57BL/6N, C57BL/6J, FVB/N and SWISS mice. Frozen zygotes may be purchased from commercial vendors (e.g., Kit Eazygote from Janvier Labs).

Instead of Cas9 other CRISPR nucleases like Cas12a can be used as well. For Cas12a use the same concentration of reagents in the electroporation mix as described for Cas9. Cas12a does not require gRNA annealing as it works without a tracrRNA. It is advisable to include the IDT electroporation enhancer at 4 µM (i.e., 0.8 µl of 100 µM in 20 µl of total mix) and to use the mix immediately after preparation for the electroporation as Cas12a can have indiscriminate ssDNA nuclease activity. For the same reason the ssODN should be stabilized with with PS (Phosphorothioate) bonds at each end (the last 3 nucleotide bonds are sufficient).

References:

- Behringer, R., et al., (2014) Manipulating the mouse embryo : a laboratory manual, Fourth edition. ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York
- Chu, V.T., et al., (2016) Efficient generation of Rosa26 knock-in mice using CRISPR/Cas9 in C57BL/6 zygotes. BMC Biotechnol 16, 4.



Materials

MATERIALS


- ⊗ Alt-R® S.p. Cas9 Nuclease V3 **Integrated DNA Technologies, Inc. (IDT) Catalog #1081058**
- ⊗ Alt-R® A.s. Cas12a Ultra Nuclease **Integrated DNA Technologies, Inc. (IDT) Catalog #10001272**
- ⊗ Alt-R® CRISPR-Cas9 tracrRNA **Integrated DNA Technologies, Inc. (IDT) Catalog #1072532**
- ⊗ Alt-R® CRISPR-Cas9 crRNA **Integrated DNA Technologies, Inc. (IDT)**
- ⊗ ssODN (Ultramer DNA Oligonucleotides) **Integrated DNA Technologies, Inc. (IDT)**
- ⊗ Alt-R® Cas12a Electroporation Enhancer **Integrated DNA Technologies, Inc. (IDT) Catalog #1076300**
- ⊗ Opti-MEM™ Reduced Serum Medium **Thermo Fisher Scientific Catalog #31985062**
- ⊗ M2 medium **Merck MilliporeSigma (Sigma-Aldrich) Catalog #M7167**
- ⊗ KSOM **Merck MilliporeSigma (Sigma-Aldrich) Catalog #MR-106-D**
- ⊗ M16 Medium
- ⊗ T10E0.1 Buffer
- ⊗ Electroporation Cuvette 1mm **Bio-Rad Laboratories Catalog #1652089**
- ⊗ 60 mm Center Well Organ Culture Dish **Corning Catalog #353037**

Troubleshooting

guide RNA annealing

- 1 Resuspend lyophilized crRNA, tracrRNA and ssODN in T₁₀E_{0.1} buffer to 100 µM (e.g., 5 nmol in 50 µl)
(Store at -80 °C until use)
- 2 Combine 5 µl crRNA (100 µM) and 5 µl tracrRNA (100 µM) in a nuclease-free PCR tube to yield an equimolar crRNA:tracrRNA duplex solution of 50 µM
- 3 Heat to 95 °C for 5 min and cool down at 5 °C/ min in a thermocycler
(crRNA:tracrRNA duplex can be stored for months at -80 °C)

Preparation of the electroporation mix

- 4 Add 1.6 µl crRNA:tracrRNA duplex and 1.3 µl Cas9 nuclease to 15.1 µl Opti-MEM in a nuclease-free tube and vortex
(Instead of Cas9 other CRISPR nucleases like Cas12a can be used as well. See Guidelines & Warnings)
- 5 Incubate mix at room temperature for 10 min
 00:10:00
- 6 Place tube on ice, add 2 µl ssODN and vortex
(The ssODN may be left out if desired but must subsequently be compensated by 2 µl Opti-MEM to reach a total electroporation mix of 20 µl)
- 7 Quick-spin at 4 °C and keep tube on ice until use

- 8 Summary of the 20 µl electroporation mix:

	A	B	C	D
	Reagent	Stock concentration	Final concentration	Volume
	crRNA:tracrRNA duplex	50 µM	4 µM	1.6 µl
	Cas9 nuclease	61 µM (10 µg/µl)	4 µM	1.3 µl
	ssODN (optional)	100 µM	10 µM	2.0 µl
	Opti-MEM	-	-	15.1 µl

Electroporation of zygotes

- 9 Collect zygotes from the oviducts of superovulated females as described in published protocols
- 10 Wash the zygotes in five drops of M2 medium
- 11 Wash up to 50 zygotes in one drop of Opti-MEM
(work quickly as zygotes may suffer from extended incubation in Opti-MEM)
- 12 Transfer zygotes with as little media as possible to the 20 μ l electroporation mix
(e.g., by first transferring the 20 μ l electroporation mix onto a culture dish and subsequently adding the zygotes)
- 13 Using a 100 μ l pipette set to 21 μ l volume transfer the entire drop of electroporation mix including the zygotes into a pre-warmed (37 °C) 1 mm electroporation cuvette
(Ensure retrieving all zygotes by quickly aspirating the entire drop. Slow aspiration will leave zygotes behind)
- 14 Insert the cuvette into a standard electroporator (e.g., BioRad Gene Pulser Xcell electroporator)
- 15 Apply two square wave pulses at 30 V and 3 ms duration with a 100 ms interval
- 16 Retrieve the zygotes by flushing the cuvette with 100 μ l M2 medium using a 100 μ l pipette into a culture dish (e.g. 60 mm Center Well Organ Culture Dish)
- 17 Wash the cuvette with 100 μ l M2 medium
- 18 Transfer all zygotes to a new culture dish containing pre-incubated microdrops of culture medium covered by oil. Wash the zygotes in three drops prior to culture in a fourth drop of culture medium.
(Alternatively, a dish with 500 μ l pre-incubated culture medium without oil may be used)
- 19 Incubate zygotes in an incubator until the two-cell stage and transfer the developed embryos into pseudopregnant foster mice
(Embryos may also be transferred at the one-cell stage)