VERSION 1

APR 17, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.7kzhkx6

Protocol Citation: Eric Danner 2020. Simple Step to Increase Gibson Assembly Efficiency . **protocols.io** <https://dx.doi.org/10.17504/protocols.io.7kzhkx6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Sep 23, 2019

Last Modified: Apr 17, 2020

PROTOCOL integer ID:
 28025

Simple Step to Increase Gibson Assembly Efficiency V.1

Eric Danner¹

¹Max Delbrück Center

Eric Danner
 Max Delbrück Center

ABSTRACT

The Gibson assembly solution is extremely toxic to some bacteria cells.

Doing a solution change after assembly and before transforming the bacteria will significantly improve bacteria survival. This increases 10-100x the amount of colonies.

MATERIALS

MATERIALS

Gibson Assembly Master Mix - 50 rxns **New England Biolabs Catalog**
#E2611L

NucleoSpin® Gel and PCR Clean-up **Macherey and Nagel Catalog**
#740609.10

Do Normal Assembly

- 1** Do a normal Gibson assembly. Incubate in the PCR machine 60 min @ 50C
- 2** Remove the assemblies. Do a PCR quickchange/ gel extract column. This will remove the Gibson Assembly solution but keep the 100ng (or however much) of DNA.

Elute in 15u -20u TE or water
- 3** Take the 15-20ul assembled plasmid and heat shock your bacteria. 30 second heat shock. 2 min ice. Add 200ul media and shake for 45 min. Carb can be plated immediately.

Plate