

Sep 05, 2025

Version 3

Simulation and Flux Analysis of β -Oxidation Using MATLAB: A Stepwise ODE-Based Protocol V.3

DOI

<https://dx.doi.org/10.17504/protocols.io.x54v95yepl3e/v3>

Ethan Shaw¹, Henry Chuo²

¹University of Houston, Baylor College of Medicine; ²University of Houston



Ethan Shaw

Baylor College of Medicine

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.x54v95yepl3e/v3>

Protocol Citation: Ethan Shaw, Henry Chuo 2025. Simulation and Flux Analysis of β -Oxidation Using MATLAB: A Stepwise ODE-Based Protocol. protocols.io <https://dx.doi.org/10.17504/protocols.io.x54v95yepl3e/v3> Version created by [Ethan Shaw](#)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: September 04, 2025

Last Modified: September 05, 2025

Protocol Integer ID: 226521

Keywords: oxidation, using ode45, using matlab, reaction kinetic, matlab, defining reaction kinetic

Abstract

This protocol outlines the complete process of modeling mitochondrial β -oxidation as a system of first-order differential equations, beginning with stoichiometric derivation and matrix reduction, and culminating in a MATLAB-based simulation. The user is guided through each step of linearizing the pathway, defining reaction kinetics, and numerically solving the system using ode45.

Materials

MATLAB (R2020b or later) — Required for numerical integration of ODE systems, matrix computation, and time-series visualization. The protocol uses built-in functions such as ode45, linspace, and plot, and is compatible with both MATLAB desktop and MATLAB Online.

Pathway Linearization and Stoichiometric Modeling

- 1 Construct a linearized kinetic representation of your metabolic pathway by referencing an established biochemical model. Then, decompose the pathway into a discrete, stepwise system of reactions in which each metabolite is uniquely identified, each enzymatic transformation is assigned a distinct reaction velocity (V_n), and each associated parameter (e.g., cofactors, substrates, rate constants) is explicitly mapped.
- 1.1 In Step 2, the biochemical β -oxidation pathway is outlined using conventional nomenclature, showing the conversion of long-chain fatty acids into acetyl-CoA through sequential enzymatic reactions, along with the involvement of key cofactors (ATP, FAD, NAD^+ , CoA-SH). Each intermediate and catalytic step is named and associated with its corresponding transformation.
- 1.2 In Step 3, the pathway is converted into a modular kinetic model, where each metabolite is denoted by a letter (A through F), and each enzymatic reaction is represented by a velocity term (*V1 through V10*). Cofactors (K through L) are similarly identified. This form isolates the stoichiometric and kinetic relationships needed for constructing ordinary differential equations (ODEs) for dynamic simulation and parameter analysis.
- 2

Fatty Acid β -Oxidation Pathway

Long Chain Fatty Acid

Thiokinase, Mg^{2+}

$ATP \rightarrow AMP + PP_i, CoA-SH$



Fatty acyl-CoA

Acyl-CoA dehydrogenase

(Dehydrogenation)

$FAD \rightarrow FADH_2$



trans- Δ^2 -Enoyl-CoA

Enoyl-CoA hydratase

(Hydration)

H_2O



β -Hydroxyacyl-CoA

β -hydroxyacyl-CoA dehydrogenase

(Dehydrogenation)

$NAD^+ \rightarrow NADH + H^+$



β -Ketoacyl-CoA

Acyl-CoA acetyltransferase


(Cleavage)

$CoA-SH$



Fatty acyl-CoA (14C) + Acetyl-CoA (2C)

3



4 Derive a set of first-order differential equations from the multi-step biochemical pathway in Step 3. Write out the reaction in the pathway, specifying reactants and products along with their stoichiometric coefficient. A negative coefficient indicates the substrate is a reactant, and a positive coefficient indicates the substrate is a product of the reaction.

5

$$\frac{dA}{dt} = V_1 - V_2$$

$$\frac{dB}{dt} = V_2 - V_3$$

$$\frac{dC}{dt} = V_3 - V_4$$

$$\frac{dD}{dt} = V_4 - V_5 - V_6$$

$$\frac{dE}{dt} = V_5 - V_7$$

$$\frac{dF}{dt} = V_6 - V_{10}$$

$$\frac{dG}{dt} = -V_8$$

$$\frac{dH}{dt} = V_8$$

$$\frac{dI}{dt} = V_9$$

$$\frac{dJ}{dt} = -V_9$$

$$\frac{dK}{dt} = -V_{10}$$

$$\frac{dL}{dt} = V_{10}$$

$$\frac{dA}{dt} = V_1 - V_2$$

$$\frac{dB}{dt} = V_2 - V_3$$

$$\frac{dC}{dt} = V_3 - V_4$$

$$\frac{dD}{dt} = V_4 - V_5 - V_6$$

$$\frac{dE}{dt} = V_5 - V_7$$

$$\frac{dF}{dt} = V_6 - V_{10}$$

$$\frac{dG}{dt} = -V_8$$

$$\frac{dH}{dt} = V_8$$

$$\frac{dI}{dt} = V_9$$

$$\frac{dJ}{dt} = -V_9$$

$$\frac{dK}{dt} = -V_{10}$$

$$\frac{dL}{dt} = V_{10}$$

6 Create a matrix corresponding to the first order differential equations, where each row represents a substrate, and each column represents a reaction in the pathway. Fill in entries of matrix with the corresponding stoichiometric coefficients determined in the previous step.

7

$$\begin{pmatrix} & V_1 & V_2 & V_3 & V_4 & V_5 & V_6 & V_7 & V_8 & V_9 & V_{10} \\ A & 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ B & 0 & 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ C & 0 & 0 & 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ D & 0 & 0 & 0 & 1 & -1 & -1 & 0 & 0 & 0 & 0 \\ E & 0 & 0 & 0 & 0 & 1 & 0 & -1 & 0 & 0 & 0 \\ F & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ G & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ H & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ J & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ K & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ L & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

- 8 Transform the stoichiometric matrix to its row-reduced echelon form by performing Gaussian elimination with back substitution. Using this, we can define dimensions, rank, and nullity of the matrix. We can also define the number of free variables in the matrix.

9

$$\begin{pmatrix} & V_1 & V_2 & V_3 & V_4 & V_5 & V_6 & V_7 & V_8 & V_9 & V_{10} \\ A & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ B & 0 & 1 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ C & 0 & 0 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ D & 0 & 0 & 0 & 1 & 0 & 0 & -1 & 0 & 0 & 0 \\ E & 0 & 0 & 0 & 0 & 1 & 0 & -1 & 0 & 0 & 0 \\ F & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ G & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ H & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ J & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ K & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ L & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} V_1 \\ V_2 \\ V_3 \\ V_4 \\ V_5 \\ V_6 \\ V_7 \\ V_8 \\ V_9 \\ V_{10} \end{pmatrix} = 0$$

Row Reduced Echelon Form of the stoichiometric matrix for the β -oxidation pathway.

10

Dimensions = 10, Rank = 9, Nullity = 1, 1 Free Variable at V_7

11 Convert the Row Reduced Echelon Form into a system of equations with each equation corresponding to a substrate row. Make each of the equations equal to 0, then use algebraic manipulation to make an equation for each reaction (V_1 , V_2 , V_3 , etc.).

12



$$V_1 - V_7 = 0$$

$$V_2 - V_7 = 0$$

$$V_3 - V_7 = 0$$

$$V_4 - V_7 = 0$$

$$V_5 - V_7 = 0$$

$$V_6 = 0$$

$$V_7$$

$$V_8 = 0$$

$$V_9 = 0$$

$$V_{10} = 0$$



$$V_1 = V_7$$

$$V_2 = V_7$$

$$V_3 = V_7$$

$$V_4 = V_7$$

$$V_5 = V_7$$

$$V_6$$

$$V_7 = V_7$$

$$V_8$$

$$V_9$$

$$V_{10}$$

13

$$J = \alpha_1 K_1 \begin{pmatrix} V_1 \\ V_2 \\ V_3 \\ V_4 \\ V_5 \\ V_6 \\ V_7 \\ V_8 \\ V_9 \\ V_{10} \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad V_7 \quad K_1$$

MATLAB Implementation and Dynamic Simulation

- 14 Write a new MATLAB function titled ode_system.m to define the ODE system governing β -oxidation dynamics. See Step 15 for confirmation.
- 14.1 At the top of the file, unpack the state vector y into individual metabolite variables (a through l) representing concentrations of each species.
- 14.2 Below, define all kinetic rate constants (k1 through k5) with realistic values derived from literature or database sources, keeping units in 1/s or 1/mM·s.
- 14.3 Define the full set of reaction velocities (V1–V10), ensuring correct substrate and cofactor dependencies for each enzymatic step. Note that V1 and V10 are equivalent.

- 14.4 Express each ODE as a difference in velocities, such that the change in each metabolite is written in the form $dx_{dt} = V_i - V_j$, avoiding invalid terms such as $+V_{10}$ or $+V_7$ in the final equations.

15

```
%% Establishing K-Vals, Reaction-V., and ODEs
function dydt = ode_system(~, y)
% Unpack variables
A = y(1); B = y(2); C = y(3); D = y(4); E = y(5);
F = y(6); G = y(7); H = y(8); I = y(9); J = y(10);
K = y(11); L = y(12);

%% Define reaction rate constants (realistic estimates in 1/s or 1/mM.s)
% [REPLACE IF/WHEN POSSIBLE]
k1 = 0.05; %Thiokinase [CORRECT]
k2 = 0.9; % Acyl-CoA dehydrogenase reaction rate [CORRECT]
%k2 = 0.01; % Modified k2 value to simulate MCADD -- ONLY UNCOMMENT IF SIMULATING MCADD
k3 = 1.2; % Enoyl-CoA hydratase reaction rate [CORRECT]
k4 = 1.0; %  $\beta$ -Hydroxy acyl-CoA dehydrogenase reaction rate [CORRECT]
k5 = 1.1; % Thiolase reaction rate [CORRECT] -- Thiolase is the shorthand for the Acyl-CoA Acetyltransferase Cleavage

%% Define reaction velocities [CORRECT]
%V1 and V10 are actually the exact same
V1 = k1 * K;
V2 = k2 * A * G;
V3 = k3 * B;
V4 = k4 * C * I;
V5 = k5 * D;
V6 = k5 * D;
V7 = k2 * E;
V8 = k2 * A * G;
V9 = k4 * C * I;
V10 = k1 * K;

%% Define ODEs [CORRECT]
% NO +V10 (dF_dt), or +V7 (dE_dt)
dA_dt = V1 - V2;
dB_dt = V2 - V3;
dC_dt = V3 - V4;
dD_dt = V4 - V6 - V5;
dE_dt = V5 - V7;
dF_dt = V6;
dG_dt = -V8;
dH_dt = V8;
dI_dt = V9;
dJ_dt = -V9;
dK_dt = -V10;
dL_dt = V10;
%Sample Corrective ODE
%dz_dt = V10 - V1 + V7;

%% Return derivative values
dydt = [dA_dt; dB_dt; dC_dt; dD_dt; dE_dt; dF_dt;
        dG_dt; dH_dt; dI_dt; dJ_dt; dK_dt; dL_dt];

end
```

- 16 In your main MATLAB script, define initial conditions for all 12 metabolites as scalar values in millimolar (mM), referencing physiologically relevant ranges for fatty acid oxidation. See Step 17 for confirmation.
- 16.1 Specify a time vector using `linspace(0,10,100)` to define a uniform time domain from 0 to 10 seconds, with 100 stored time points.
- 16.2 Call the MATLAB solver using `[t, y] = ode45(@ode_system, tspan, [A0 B0 C0 D0 E0 F0 G0 H0 I0 J0 K0 L0]);` to simulate the ODE system and record the time-course evolution of each metabolite.

17

```
%% Define realistic initial conditions (in mM)
A0 = 0.5; % Fatty Acyl-CoA ~ 0.5 mM (depends on lipid metabolism rate)
B0 = 0.05; % Trans  $\Delta^2$  Enoyl-CoA ~ 0.05 mM (intermediate product)
C0 = 0.03; %  $\beta$ -Hydroxy acyl-CoA ~ 0.03 mM (intermediate product)
D0 = 0.02; %  $\beta$ -Keto acyl-CoA ~ 0.02 mM (intermediate product)
E0 = 1.5; % Acetyl-CoA ~ 1.5 mM (enters citric acid cycle)
F0 = 0.2; % Shortened Acyl-CoA ~ 0.2 mM (for next cycle)
G0 = 0.1; % FAD ~ 0.1 mM (electron carrier)
H0 = 0.05; % FADH2 ~ 0.05 mM (reduced electron carrier)
I0 = 1.0; % NAD+ ~ 1.0 mM (electron acceptor for oxidation)
J0 = 0.5; % NADH ~ 0.5 mM (reduced electron carrier)
K0 = 2.0; % ATP ~ 2.0 mM (energy donor for fatty acid activation)
L0 = 0.1; % AMP ~ 0.1 mM (byproduct of ATP hydrolysis in activation step)

%% Figure Specification
%% Define time span with stored time values
tspan = linspace(0, 10, 100); % Store 100 evenly spaced time points from 0 to 10

%% Solve ODEs using ode45
[t, y] = ode45(@ode_system, tspan, [A0 B0 C0 D0 E0 F0 G0 H0 I0 J0 K0 L0]);
```

18



beta_oxidation_model.m 22.6KB

For further features such as custom figure generation, reaction logging, or MCADD simulation toggles, download the complete model file: *betaoxidationmodel.m* – which contains embedded comments and configurable options for simulation refinement.