VERSION 1

NOV 02, 2018

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.u9yez7w

Protocol Citation: Samantha A Morris, Brent A. Bidy, Wenjun Kong, Kenji Kamimoto, Chuner Guo, Sarah Waye, Tao Sun 2018. Single-cell mapping of lineage and identity via CellTagging.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.u9yez7w>

MANUSCRIPT CITATION:

<https://www.biorxiv.org/content/early/2017/04/28/127860>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development

We are still developing and optimizing this protocol

Created: Nov 02, 2018

Single-cell mapping of lineage and identity via CellTagging V.1

Samantha A Morris¹, Brent A. Bidy¹, Wenjun Kong¹, Kenji Kamimoto¹, Chuner Guo¹, Sarah Waye¹, Tao Sun¹

¹Washington University, Saint Louis

Morris Lab

Samantha A Morris

Washington University, Saint Louis

ABSTRACT

Single-cell technologies are offering unprecedented insight into complex biology, revealing the behavior of rare cell populations that are typically masked in bulk population analyses. One current limitation of single-cell approaches is that lineage relationships are lost as a result of cell processing, restricting interpretations of the data collected. Elegant computational approaches have been developed in an effort to infer these missing observations, but it remains a challenge to reconstruct true reprogramming trajectories using these tools. Although sophisticated lineage tracing solutions to connect cell history with fate are emerging, these protocols are either not compatible with high-throughput scRNA-seq, or require genome editing strategies that are not readily deployed in some systems. Here, our protocol describes a single-cell resolution clonal tracking approach, 'CellTagging', based on combinatorial cell indexing, permitting the parallel capture of lineage information and cell identity. CellTagging integrates with high-throughput single-cell RNA-sequencing, where iterative rounds of cell labeling enable the construction of multi-level lineage trees. This straightforward lentiviral-labeling approach can be applied to an array of cell biological applications to simultaneously profile lineage and identity, at single-cell resolution.

MATERIALS

MATERIALS

⊗ Stellar Competent Cells **Takarabio Catalog #636763**

⊗ CellTag pooled library V1 **addgene Catalog #115643**

⊗ CellTag pooled library V2 **addgene Catalog #115644**

⊗ CellTag pooled library V3 **addgene Catalog #115645**

PROTOCOL integer ID:
17432

Keywords: cell barcoding,
lineage tracing, single-cell
analysis, direct lineage
reprogramming

For generation of lentivirus, the appropriate institutional regulations must be satisfied.

Amplification of CellTag library

1

Thaw Stellar Competent Cells in an ice bath just before use.

2

After thawing, mix 100 uL of cells with 10-50 ng of CellTag V1 DNA in a 1.5-mL microcentrifuge tube.

3

Place transformation mixture on ice for 30 minutes.

4

Heat shock the cells for exactly 60 seconds at 42 °C .

5

Place tube on ice for 1-2 minutes.

6

Add SOC medium to bring the final volume to 1000 uL.

7

Incubate by shaking (~250 rpm) for 1 hour at 37 °C .

- 8 Take 6 μL of the recovery. Use 1 μL of this to prepare a 1:10 and then a 1:100 dilution. Plate both dilutions and the rest of the 5 μL of recovery onto three separate LB/Amp plates. Grow overnight at 37 °C .
- 9 Add the rest of the recovery to 500 mL of LB+Amp. Grow overnight while shaking (~250 rpm) at 37 °C .
- 10 After counting the CFUs from the LB/Amp plates the next day, calculate the transformation efficiency.
- 11 Harvest the cells from the liquid culture and use multiple Maxiprep columns (Sigma or Zymogen) to purify the library.
- 12