JUN 27, 2017

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.invcde6

Protocol Citation: Gloria Patricia Barrera¹, Laura Fernanda Villamizar⁴, Carlos Espinel¹, Edgar Mauricio Quintero², Mariano Nicolás Belaich³, Deisy Liseth Toloza¹, Pablo Daniel Ghiringhelli^{3*}, German Vargas² 2017. Single-Strand Conformation Polymorphism (SSCP) for *Diatraea* species identification. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.invcde6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Jun 27, 2017

Last Modified: Mar 25, 2018

PROTOCOL integer ID:
 6581

Single-Strand Conformation Polymorphism (SSCP) for *Diatraea* species identification

Gloria Patricia Barrera¹, Laura Fernanda Villamizar⁴, Carlos Espinel¹, Edgar Mauricio Quintero², Mariano Nicolás Belaich³, Deisy Liseth Toloza¹, Pablo Daniel Ghiringhelli^{3*}, German Vargas²

¹Centro de investigación Tibaitatá, Corporación Colombiana de Investigación Agropecuaria CORPOICA, Km 14 Vía Mosquera 250047, Cundinamarca, Colombia ²Centro de investigación de la caña de azúcar de Colombia (CENICAÑA). Calle 58 norte No. 3BN-110 Cali – Colombia. ³Laboratorio de Ingeniería Genética y Biología Celular y Molecular—Área Virosis de Insectos (LIGBCM—AVI), Dto. de Ciencia y Tecnología, Universidad Nacional de Quilmes, Roque Saenz Peña 352, Bernal, Provincia de Buenos Aires, 1876, Argentina. ⁴AgResearch Ltd. Lincoln Research Centre. Christchurch 8140, New Zealand

Gloria Barrera

ABSTRACT

We propose a Single-Strand Conformation Polymorphism (SSCP) methodology to recognize each species as an effective tool in *Diatraea* species identification.

MATERIALS

STEP MATERIALS

☒ DNeasy Blood & Tissue Kit, QIAGEN Contributed by users Catalog #Cat No./ID: 69504

☒ Taq polymerase Promega Catalog #M3001

☒ SYBR Thermo Fisher Scientific Catalog #S33102

☒ DNeasy Blood & Tissue Kit, QIAGEN Contributed by users Catalog #Cat No./ID: 69504

☒ Taq polymerase Promega Catalog #M3001

☒ SYBR Thermo Fisher Scientific Catalog #S33102

- 1 Dissection and DNA extraction:** Abdomens from insects were dissected and maintained in ethanol at 70 % v/v in sterile distilled water to DNA extractions. DNA of male insects were obtained of complete abdomen by using commercial kit (DNeasy Blood & Tissue Kit, QIAGEN) and quantified by spectrophotometry (Nanodrop 1000, Thermo-Fisher).

✂ DNeasy Blood & Tissue Kit, QIAGEN **Contributed by users Catalog #Cat No./ID: 69504**

- 2 COII amplification:** In vitro amplifications of CO II fragments were carried out by the Polymerase Chain Reaction method (PCR) in standard conditions (final volume 25 µL) using Taq polymerase (PROMEGA), 50 ng of template and the primers previously described and used in Diatraea: A-298 (5'-ATTGGACATCAATGATATTGA-3') and B-tLYS (5' GTTTAAGAGACCACTACTTG-3') (Simon et al. 1994; Lange et al. 2004, Palacio et al. 2010). Thermal cycling was performed using the following conditions: 1 cycle at 95 °C for 3 min; 34 cycles at 95 °C for 10 s; 53 °C for 45 s, and 72 °C for 30 s; and one cycle at 72 °C for 5 min. Aliquots of amplification products were resolved in 1% w/v agarose gel electrophoresis and later stained with SYBR-Safe (Invitrogen). Amplicons were used for SSCP methodology.

✂ Taq polymerase **Promega Catalog #M3001**

✂ SYBR **Thermo Fisher Scientific Catalog #S33102**

- 3 Single-Strand Conformation Polymorphism (SSCP) methodology:** The amplicons were prepared for SSCPs. To this, 1 µL of PCR products amplified using A-298/B-tLYS were mixed with 9 µL of sample buffer (95% v/v Formamide, 20mM EDTA, 0.05% w/v bromophenol blue and 0.05% w/v Xylene-cyanol, in distilled water), denatured at 95 °C for 5 min and immediately stored in ice bath. Then, 1.5 µL of all samples were loaded in 6% non-denaturing polyacrylamide gel (49 acrylamide: 1 bis-acrylamide), and separated by electrophoresis for 4 hours at 600V (BioRad Sequi-Gen GT Nucleic Acid Electrophoresis Cell, 38 x 50 cm). Later, PAGEs were stained using and adapted silver nitrate method. Briefly, gels were fixed for 3 min (10% v/v ethanol, 1% v/v acetic acid in distilled water), oxidized for 3 min (1.5 % v/v nitric acid in distilled water), stained for 20 min (0.1% w/v silver nitrate, 0.045 % v/v formaldehyde in distilled water), revealed for approximately 5 min (3% w/v NaCO₃, 0.02 % v/v formaldehyde in distilled water), and the reactions were stopped with an acetic acid solution (5 % v/v in distilled water). Results were documented by digital photography.