APR 19, 2018

Soil microbial community structure -PLFAs

PLOS One

Jing Li¹

¹Chinese Academy of Agricultural Sciences

Jing Li

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.nggdbtw

External link:

<https://doi.org/10.1371/journal.pone.0195144>

Protocol Citation: Jing Li

2018. Soil microbial community structure -PLFAs.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.nggdbtw>

MANUSCRIPT CITATION:

Li J, Wu X, Gebremikael MT, Wu H, Cai D, Wang B, Li B, Zhang J, Li Y, Xi J (2018) Response of soil organic carbon fractions, microbial community composition and carbon mineralization to high-input fertilizer practices under an intensive agricultural system. PLoS ONE 13(4): e0195144. doi: [10.1371/journal.pone.0195144](https://doi.org/10.1371/journal.pone.0195144)

[4](https://doi.org/10.1371/journal.pone.0195144)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Feb 26, 2018

Last Modified: Apr 19, 2018

PROTOCOL integer ID:
10472

- 1 Three grams of freeze-dried soil samples were used to extract the lipids by a single-phase CHCl_3 : methanol: citrate buffer (15.2 mL at a 1:2:0.8 volume ratio).
- 2 The soil extracts were fractionated into neutral lipids, glycolipids, and polar lipids using a silica-bonded phase column (SPE-Si, Supelco, Poole, UK) with CHCl_3 , acetone and methanol, respectively.
- 3 The recovered polar lipids were saponified and methylated to fatty acid methyl esters (FAME).
- 4 FAMES were quantified by a gas chromatograph (N6890, Agilent) and identified with an MIDI Sherlock Microbial Identification System (Version 4.5, MIDI, Inc., Newark, DE).
- 5 We divided all PLFAs into 6 microbial groups based on previously published PLFA biomarker data: Gram-positive (G+) bacteria (i14:0, i15:0, a15:0, i16:0, i17:0, a17:0), Gram-negative (G-) bacteria (16:1 ω 7c, 16:1 ω 9c, 17:1 ω 8c, 18:1 ω 5c, 18:1 ω 7c, cy17:0, cy19:0), actinomycetes (10Me16:0, 10Me17:0, 10Me18:0), saprophytic fungi (18:1 ω 9c and 18:2 ω 6c) and AMF (16:1 ω 5c).