VERSION 2

JUL 23, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.biahkab6

Document Citation: Marco Cosentino, Elisa Storelli, Alessandra Luini, Massimiliano LM Legnaro, Emanuela Rasini, Marco Ferrari, Franca Marino 2020. SOLUTION- 02 - Phosphate Buffered Saline (PBS). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.biahkab6>

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Jul 07, 2020

Last Modified: Jul 27, 2020

DOCUMENT integer ID:
 38953

🌐 SOLUTION- 02 - Phosphate Buffered Saline (PBS) V.2

Marco Cosentino¹, Elisa Storelli¹, Alessandra Luini¹, Massimiliano LM Legnaro¹, Emanuela Rasini¹, Marco Ferrari¹, Franca Marino¹

¹Center for Research in Medical Pharmacology, University of Insubria (Varese, Italy)

Elisa Storelli

Center for Research in Medical Pharmacology, University of I...

ABSTRACT

This recipe is used in the following protocols:

- Separation and purification of human PBMC from FRESH BLOOD
- Separation and purification of human PBMC from BUFFY COAT
- Magnetic bead-based CD4+ T cell isolation from PBMCs with Dynabeads: CD4 Positive Isolation Kit
- Magnetic bead-based TREG-TEFF cell isolation from PBMC with Miltenyi CD4+CD25+ Regulatory T cell Isolation Kit
- Staining of human PBMC or ISOLATED SUBSETS with Cell Proliferation Dye-eFluor™ 670 (CPD-eFluor670) for cell proliferation evaluation by Flow Cytometry

Phosphate-buffered saline (PSB 1X)

NaCl - 8 g (137 mM)

KCl - 0.2 g (2.7 mM)

Na₂HPO₄ - 1.44 g (10 mM)

KH₂PO₄ - 0.24 g (1.8 mM)

To prepare 1 L of PBS 1X, dissolve the reagents listed above in 800 mL of ultrapure H₂O. Adjust the pH to 7.4 with HCl or NaOH, and then add H₂O to 1 L.

NaCl code: S9625, Sigma

KCl code: P9541, Sigma

Na₂HPO₄ code: 1.06585, Sigma

KH₂PO₄ code: P0662, Sigma

Fonte: (Cold Spring Harbor Laboratory Press 2006)