

Jul 23, 2020

SOLUTION- 08 - Türk solution

DOI

dx.doi.org/10.17504/protocols.io.biwvkfe6

Marco Cosentino¹, Elisa Storelli¹, Alessandra Luini¹, Massimiliano LM Legnaro¹, Emanuela Rasini¹, Marco Ferrari¹, Franca Marino¹

¹Center for Research in Medical Pharmacology, University of Insubria (Varese, Italy)



Farmacologia Medica

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.biwvkfe6

Document Citation: Marco Cosentino, Elisa Storelli, Alessandra Luini, Massimiliano LM Legnaro, Emanuela Rasini, Marco Ferrari, Franca Marino 2020. SOLUTION- 08 - Türk solution. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.biwvkfe6>

License: This is an open access document distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: July 23, 2020

Last Modified: November 21, 2020

Document Integer ID: 39605



Abstract

This recipe is used in the following protocols:

- **PBMC- 01a** - Isolation of Human PBMC from Buffy Coat
- **PBMC- 01b** - Isolation of Human PBMC from Whole Blood
- **PBMC- 02** - CD4+ T cell Isolation from PBMC with "Dynabeads CD4 Positive Isolation Kit"
- **PBMC- 03** - TEFF+TREG Isolation from PBMC with "Miltenyi CD4+CD25+ Regulatory T cell Isolation Kit"
- **PMN- 01a** - Isolation of Human PMN from Buffy Coat
- **PMN- 01b** - Isolation of Human PMN from Whole Blood



Türk solution composition (v/v):

- Gentian Violet 0.02%
- Acetic Acid 3% prepared in H₂O.

Storage: 4°C in the dark covered with aluminum foil (**photosensitive**)

Gentian violet: Marco Viti

Acetic acid catalog number: A6283 Sigma Italy